

TABLE 2. Percentage distribution of women aged 18–49 at risk of pregnancy, by the effectiveness of their primary contraceptive method, according to jurisdiction, 2019†

Jurisdiction	Highly effective (permanent‡)	Highly effective (LARC§)	Moderately effective**	Least effective††	Unspecified method	No method	Total
Alabama	22.4	6.5	16.9	24.8	4.5	25.0	100
Arizona	21.8	8.8	16.6	19.2	4.6	29.1	100
Arkansas	19.3	6.9	17.7	21.1	4.4	30.6	100
Connecticut	13.2	11.8	17.6	22.9	6.0	28.5	100
Delaware	14.0	9.2	20.7	22.2	5.6	28.2	100
Florida	18.4	8.9	17.8	20.0	5.8	29.2	100
Georgia	14.3	8.4	19.1	19.8	4.3	34.2	100
Hawaii	11.1	11.8	15.0	15.6	6.2	40.3	100
Idaho	26.2	17.7	13.5	14.5	4.7	23.3	100
Illinois	15.2	7.5	17.2	25.6	5.7	28.7	100
Indiana	19.1	9.4	16.6	20.0	4.1	30.8	100
Iowa	24.9	14.3	18.5	16.2	3.3	22.8	100
Kansas	19.3	13.1	19.0	16.2	4.9	27.4	100
Louisiana	16.7	8.2	18.5	25.2	3.1	28.3	100
Maryland	14.4	11.4	17.4	23.7	6.1	27.0	100
Massachusetts	12.5	14.7	21.5	23.9	5.2	22.1	100
Minnesota	18.2	15.1	19.7	16.2	4.9	25.9	100
Mississippi	23.7	6.9	16.6	22.9	1.7	28.2	100
Missouri	23.8	10.5	17.6	20.2	4.4	23.6	100
Montana	25.9	18.8	14.9	15.4	3.8	21.1	100
Nebraska	18.0	12.5	19.7	21.0	2.8	26.1	100
New Mexico	22.5	15.0	12.1	18.6	‡‡	29.0	100
New York	7.2	11.1	14.2	26.5	6.9	34.1	100
North Carolina	17.8	13.6	16.6	16.1	3.3	32.6	100
Ohio	19.4	10.4	20.1	15.4	4.5	30.2	100
Oklahoma	25.1	9.9	17.9	16.1	5.9	25.2	100
Oregon	21.8	20.6	16.2	16.7	4.3	20.4	100
Pennsylvania	19.5	9.3	20.6	19.8	‡‡	27.1	100
Puerto Rico	36.8	3.9	7.8	18.9	3.6	29.0	100
Rhode Island	13.4	12.3	23.0	19.7	4.8	26.8	100
South Carolina	16.6	8.3	19.7	21.3	4.1	30.1	100
South Dakota	22.7	12.0	18.7	18.1	3.4	25.1	100
Tennessee	21.8	7.1	22.2	19.1	4.3	25.5	100
Utah	21.3	24.8	15.0	14.6	3.3	21.0	100
Virginia	16.0	12.1	19.1	17.5	5.8	29.5	100
West Virginia	29.1	10.0	15.6	16.3	4.6	24.5	100
Wisconsin	17.9	13.6	23.9	19.0	‡‡	24.4	100
Wyoming	22.7	11.8	14.7	17.6	6.6	26.7	100

†Women at risk of pregnancy are those aged 18–49 who are sexually active with one or more male partners, are not currently pregnant or postpartum, and have not had a hysterectomy. ‡Female sterilization or male sterilization, as reported by respondent. §IUDs and contraceptive implants. **Injectables, pills, patches and vaginal rings. ††Male and female condoms, diaphragm, cervical cap, sponge, rhythm method, natural family planning, withdrawal, spermicidal foam/jelly/film/cream and emergency contraception. ‡‡Estimate was suppressed (has a denominator of fewer than 50 respondents or a relative standard error greater than 30%). Note: LARC=long-acting reversible contraceptive.