TABLE 1. Percentage distribution of participants in a study of a sexual and reproductive health intervention for outof-school men, by selected demographic characteristics at baseline, according to study condition, Baltimore, 2008–2009

Characteristic	All (N=197)	Control (N=99)	Intervention (N=98)					
Mean age								
16–17	35.0	29.3	40.8					
18–19	35.5	36.4	34.7					
20–24	29.4	34.3	24.5					
Race/ethnicity								
Black	95.9	97.0	94.9					
Other	4.1	3.0	5.1					
Last grade completed								
≤9th	45.2	39.4	51.0					
10th/11th	46.7	47.5	45.9					
≥12th/GED	7.6	13.1	2.0**					
Missing	0.5	0.0	1.0					
Has health insurance								
Yes	61.4	64.6	58.2					
No	37.6	35.4	39.8					
Missing	1.0	0.0	2.0					
Visited provider in last 12 mos.†								
Yes	58.4	60.6	56.1					
No	37.1	35.4	38.8					
Missing	4.6	4.0	5.1					
Total	100.0	100.0	100.0					

**p<.01. †When sick or in need of care. *Note:* Percentages may not total 100.0 because of rounding.

TABLE 2. Selected knowledge, attitude and behavioral characteristics of participants at baseline and three-month follow-up, by study condition; and odds ratios (and 95% confidence intervals) from multivariate regression analysis assessing the likelihood of intervention participants' experiencing a change in outcome over time relative to control participants'

Characteristic	Control (N=77)		Intervention (N=75)		Odds ratio
	Baseline	Follow-up	Baseline	Follow-up	
Mean knowledge scores					
STDs (range, 0–1)	0.45 (0.27)	0.36 (0.27)**	0.46 (0.24)	0.81 (0.23)***	1.57 (1.44–1.71)***
Health care (range, 0–1)	0.36 (0.30)	0.35 (0.35)	0.35 (0.35)	0.81 (0.26)***	1.57 (1.40–1.76)***
Mean attitude scores					
No way to enjoy sex with a condom (range, 1–4)	2.74 (1.11)	2.83 (1.03)	2.93 (0.98)	3.28 (0.97)*	1.39 (0.93–2.09)
Feelings about condoms (range, 1–10)	7.97 (2.25)	7.68 (2.24)	7.78 (2.22)	8.09 (2.20)	1.62 (0.69–3.84)
Want sex with new partner, but no condom (range, 1-4)	3.05 (0.84)	3.09 (0.81)	2.92 (0.89)	3.21 (0.82)	1.20 (0.87–1.65)
Sexual behavior in last 3 mos.					
Had vaginal/oral/anal sex†	92.9	90.0	90.4	93.2	2.77 (0.46–16.53)
No.of partners†					
0	7.1	16.2	9.6	9.3	1.64 (0.47–5.70)‡
1	40.0	44.6	34.2	29.3	na
2	15.7	17.6	20.5	26.7	na
≥3	27.6	21.6*	31.1	34.7	na
Mean frequency of condom use (range, 1–4)§	2.66 (0.98)	2.44 (1.10)*	2.77 (1.05)	3.12 (1.06)*	1.83 (1.24–2.71)**
Used lubricant with condoms§	14.3	8.1	8.8	23.9*	23.61 (1.67–333.35)*
Used condom at last sex§	54.0	61.1	61.8	75.3	1.10 (0.88–1.34)
Health care use in last 3 mos.†					
Talked to provider about STDs/HIV	44.3	16.9***	26.0	45.4*	12.32 (3.81–39.79)***
Got checked for STDs/HIV	55.7	22.4***	28.8	36.0	16.62 (3.77–73.30)***

*p<.05. **p<.01.***p<.001.**Odds ratios were calculated for those who had had sex in the three months before baseline` or follow-up. ‡Assessed as a continuous variable. §Odds ratios were calculated for those who had had sex in the three months before follow-up. *Notes*: Unless otherwise noted, data are percentages. Higher scores indicate greater knowledge, more positive attitudes or greater frequency. Figures in parentheses alongside means are standard deviations. Paired t tests or McNemar's tests were used to assess differences between baseline and follow-up values. See page 36 for a description of which variables were controlled for in the different regression models. na=not applicable.