Table 2. Percentage of sexually active women who reported a bacterial STD or treatment for PID, by selected risk characteristics

Characteristic	Bacterial STD	PID
Age at first intercourse		
<15	13.0	12.1
15–16	7.0	9.9
17–18	5.3	7.8
≥19	3.4	5.6
Lifetime number of sexual partners		
1	1.4	5.0
2–3	3.2	6.8
4–5	5.9	7.3
>5	14.0	14.0
Frequency of intercourse in past 3 months†		
None	3.6	3.9
<once per week	6.9	8.7
Once per week	4.6	7.5
>once per week	6.5	9.0
Ever used a condom		
No	2.9	7.3
Yes	7.0	8.6
Used condom at last intercourse‡		
No	6.4	6.4
Yes	4.7	4.6
Ever used a condom to prevent STDs§		
No	4.5	6.8
Yes	9.9	10.7
Screened for STD in past 12 months		
No	4.6	7.2
Yes	17.7	11.6
Ever used IUD		
No	6.0	7.4
Yes	8.8	17.3
Frequency of douching		
Never	5.5	6.0
<weekly	6.3	12.1
>weekly	5.7	10.9
Had a bacterial STD		
Never	0.0	7.4
Ever	100.0	23.0

†Frequency indicated only for those reporting intercourse in the three months before the survey. ‡Based on women who reported intercourse in the three months before the survey and who used a contraceptive at last intercourse. §Includes women who reported use for both contraception and disease prevention. *Note:* Estimates are weighted.