**Table 1. Percentage distributions (and standard errors) of women aged 15–44 who are at risk of unintended pregnancy, by contraceptive use pattern during past year, according to age-group, 1995 National Survey of Family Growth**

| Age-group | Uninterrupted use | | Sporadic use | Nonuse | Total | N |
|---|---|---|---|---|---|---|
| | Effective method | Other method | | | | |
| **Total** | **61.4 (0.6)** | **29.6 (0.6)** | **5.6 (0.3)** | **3.5 (0.2)** | **100.0** | **6,828** |
| 15–17 | 23.3 (2.6)* | 48.6 (3.3)* | 20.5 (2.6)* | 7.5 (1.8)* | 100.0 | 252 |
| 18–19 | 31.3 (3.2)* | 45.5 (3.6)* | 18.4 (2.5)* | 4.8 (1.2) | 100.0 | 322 |
| 20–24 | 52.2 (1.8)* | 34.3 (1.8)* | 9.2 (1.0)* | 4.2 (0.6) | 100.0 | 1,000 |
| 25–34 | 62.4 (1.1) | 30.1 (1.1) | 3.8 (0.4) | 3.6 (0.4) | 100.0 | 2,643 |
| ≥35 | 72.9 (1.0)* | 22.6 (1.0)* | 2.2 (0.3)* | 2.4 (0.3)* | 100.0 | 2,611 |

*Significantly different from women aged 25–34 at p<.05. *Notes:* Percentages are weighted. Ns are unweighted.