

Table 3. Percentage distribution of women who obtained an abortion, by location of provider, according to residence and time period

Provider location	Total		Rural		Urban	
	1983–1984 (N=53,287*)	1993–1994 (N=53,662*)	1983–1984 (N=6,913†)	1993–1994 (N=5,913†)	1983–1984 (N=46,374‡)	1993–1994 (N=47,681‡)
Urban	93.6	94.5	66.9	83.2	97.5	95.9
Large urban	83.0	84.7	48.4	58.2	88.1	88.0
Small urban	10.6	9.8	18.5	25.0	9.4	7.9
Rural	3.5	0.3	24.8	2.7	0.3	0.0
Large rural	1.7	0.0	12.1	0.1	0.1	0.0
Small rural	1.8	0.3	12.7	2.6	0.2	0.0‡
Out of state	2.9	5.2	8.3	14.2	2.1	4.0
Idaho	0.3	0.1	2.2	0.5	0.0‡	0.0‡
Oregon	2.6	5.1	6.1	13.6	2.1	4.0
Other	0.0‡	0.0‡	0.0‡	0.1	0.0‡	0.0‡
Total	100.0	100.0	100.0	100.0	100.0	100.0

*Includes cases with missing age and county. †Includes cases with missing age; does not include cases with missing county. ‡Less than 0.05%. *Notes:* All rural-urban differences are statistically significant at $p < .01$ (by the chi-square test), except the differences in the "other" category. All interdecade differences are significant at $p < .01$ (by chi-square test), except for the "other" category and for urban residents obtaining abortions in large urban counties. Women aged 50–59 are excluded. Percentages may not add to 100 due to rounding.