Table 1. Percentage distribution of general practitioners, by opinion about who should have responsibility for a woman's decision to have an abortion, according to gestation, Great Britain, 1999

` '		
Who should have responsibility	≤12 weeks	>12 weeks

(N=702)

who should have responsibility	≤12 weeks	>12 weeks
Woman only	46.4 (42.7–50.1)	21.4 (18.3–24.4)
Woman and one doctor	7.3 (5.5–9.4)	6.0 (4.4–8.0)

oman only	46.4 (42.7–50.1)	21.4 (18.3-
oman and one doctor	7.3 (5.5–9.4)	6.0 (4.4–8
oman and two doctors	16.8 (14.0–19.6)	23.4 (20.2-
	0.4 (4.4.0.0)	20/40

Woman and two doctors	16.8 (14.0–19.6)	23.4 (20.2–26.5)
One doctor only	2.4 (1.4–3.9)	2.9 (1.8–4.4)
Two doctors only	10 5 (16 6-22 1)	36 0 (33 3-40 5)

One doctor only	2.4 (1.4–3.9)	2.9 (1.8–4.4)
Two doctors only	19.5 (16.6–22.4)	36.9 (33.3–40.5)

Two doctors only	19.5 (16.6–22.4)	36.9 (33.3-40.5)
Other/no response	7.6 (5.7–9.8)	9.5 (7.5–12.0)

Other/no response	7.6 (5.7–9.8)	9.5 (7.5–12.0)
Total	100.0	100.0

Note: Data in parentheses are 95% confidence intervals.