

**TABLE 2. Percentage distribution (and 95% confidence intervals) of currently married and sexually active unmarried young women using a modern contraceptive method, by source of most recent method, according to method**

Source	All (N=536)	Pill (N=166)	Injectable (N=239)	Implant (N=36)	Condom (N=77)
<b>Public</b>	<b>66.5</b>	<b>66.1 (57.3–73.9)</b>	<b>83.8 (76.0–89.4)</b>	<b>95.4 (83.3–98.9)</b>	<b>5.2 (1.8–13.7)</b>
Hospital	2.1	2.3 (0.8–6.2)	1.5 (0.6–3.8)	10.4 (4.4–22.5)	0.0
Health center	17.2	19.3 (13.1–27.7)	13.3 (9.2–18.9)	60.1 (41.1–76.6)	1.2 (0.2–8.1)
Health post	42.8	39.3 (30.9–48.4)	63.7 (55.9–70.9)	24.1 (11.8–43.0)	3.7 (1.0–12.5)
Other public†	4.4	5.2 (2.3–11.2)	5.3 (2.9–9.6)	0.8 (0.1–5.8)	0.2 (0.0–1.7)
<b>Private</b>	<b>26.1</b>	<b>29.9 (22.5–38.5)</b>	<b>14.6 (9.2–22.5)</b>	<b>0.0</b>	<b>61.1 (46.7–73.8)</b>
Hospital/clinic	8.2	9.1 (4.9–16.2)	11.3 (6.4–19.3)	0.0	1.0 (0.1–6.7)
Other private	17.9	20.8 (14.8–28.5)	3.3 (1.5–7.2)	0.0	60.1 (45.8–72.9)
<b>NGO/other‡</b>	<b>7.4</b>	<b>4.0 (1.6–9.7)</b>	<b>1.6 (0.6–4.1)</b>	<b>4.6 (1.1–16.7)</b>	<b>33.7 (21.5–48.5)</b>
Total	100.0	100.0	100.0	100.0	100.0

†Includes dispensaries and community health centers. ‡Includes workplace clinics, youth centers, voluntary counseling and testing centers, shops, markets and peer educators. Notes: NGO=nongovernmental organization. Data for the IUD and other methods are not presented because of small sample sizes.