Country and method	No. of episodes	All causes		Desire for	Desire for pregnancy		Method failure		Health concerns		Other	
		12	24	12	24	12	24	12	24	12	24	
MOROCCO												
All methods	2,914	35.2	53.5	9.8	18.4	12.2	20.5	11.3	18.6	7.7	12.1	
Pill	2,050	35.1	53.5	10.4	18.6	9.1	16.3	14.1	22.8	7.2	11.5	
IUD	182	17.5	34.5	1.6	5.1	4.2	9.4	8.2	17.1	4.6	8.0	
Other modern methods*	100	49.8	68.2	19.3	31.7	20.6	28.7	7.1	11.5	15.7	24.8	
Periodic abstinence	237	44.5	65.0	13.8	26.4	29.3	45.9	1.5	1.5	7.5	10.7	
Withdrawal	232	30.8	50.6	7.4	19.2	14.6	25.1	1.3	1.3	11.3	18.2	
TUNISIA												
All methods	2,289	29.5	45.6	5.8	12.3	8.4	13.0	11.5	18.7	7.5	12.2	
Pill	662	40.6	58.5	8.8	18.0	6.7	9.0	23.3	36.3	8.9	12.6	
IUD	855	147	30.2	22	6.6	22	49	8.5	16.2	24	63	
Vaginal methods	144	63.9	83.6	11.4	25.2	41.6	50.8	10.9	20.4	21.6	43.8	
Other modern methodst	133	41 4	50.8	11.3	18.0	6.5	6.5	14.1	14 1	17.6	25.2	
Periodic abstinence	300	26.1	12.6	7.2	1/ 0	14.5	24.4	0.0	0.0	67	10.8	
Withdrawal	107	20.1	42.0	1.2	0 5	14.5	24.4	0.0	0.0	15.7	10.0	
Williulawai	137	33.1	47.3	0.0	0.0	15.5	20.3	0.0	0.0	15.7	19.7	
EGYPT All methods	1 196	28.0	/8.8	53	13.8	71	14.2	12.4	21.3	67	11 0	
Dill	2,295	26.5	50.4	7.5	19.6	7.0	16.5	10.1	20.9	9.0	12.5	
	1 496	10.0	02.4	7.5	6.4	7.9	10.5	5.0	10.0	1.5	10.0	
Condom	1,400	10.9	23.0	2.3	10.4	2.0	4.1	5.0	12.1	1.5	01 5	
	200	35.1	53.4	5.3	13.4	10.4	29.4	1.3	2.5	16.9	21.5	
Other modern methods‡	128	58.7	80.0	13.7	23.2	14.0	33.7	24.7	37.7	26.1	36.4	
Breastreeding	180	11.0	56.5	0.8	14.9	5.6	21.1	0.3	5.4	4.7	31.3	
Iraditional methods§	134	41.9	63.7	5.4	22.2	30.3	41.9	1.3	1.3	10.7	19.6	
ECUADOR	4 9 4 9		40.0		10 -		17.0		45.0	10.0	40.0	
All methods	1,348	34.6	48.0	5.9	10.7	11.6	17.6	9.8	15.8	12.8	16.6	
Pill	506	37.0	55.2	7.8	14.6	6.4	13.2	18.9	29.8	9.9	13.8	
IUD	327	13.6	19.9	1.1	1.1	3.8	6.7	3.9	7.0	5.6	6.7	
Other modern methods*	177	62.4	75.8	12.8	24.3	23.6	28.7	7.1	9.6	39.2	51.1	
Periodic abstinence	245	35.3	47.9	5.1	10.1	21.5	31.2	0.8	0.8	12.2	15.0	
INDONESIA												
All methods	6,183	25.1	40.6	7.5	14.4	2.8	4.9	8.7	14.0	8.7	15.1	
Pill	2,308	26.2	43.2	9.3	18.6	2.2	3.4	10.2	15.9	6.3	14.3	
IUD	1,246	12.4	21.4	2.1	5.0	2.1	4.4	5.8	8.8	3.0	5.1	
Injectable	1,637	24.9	43.9	5.5	10.9	1.1	2.2	11.6	20.5	9.0	19.1	
Condom	272	42.0	57.5	15.8	26.9	5.7	9.4	4.0	7.8	23.9	31.9	
Periodic abstinence	190	34.9	57.3	13.7	25.6	12.2	25.8	0.9	2.3	13.4	20.8	
Withdrawal	217	46.7	61.7	21.0	36.3	8.9	16.7	1.2	1.2	25.0	26.9	
Herbs	152	41.1	60.3	18.5	33.7	7.2	12.2	6.7	7.6	17.2	26.2	
THAILAND												
All methods	3,407	36.3	50.4	14.1	23.0	3.9	5.4	6.8	10.9	17.1	23.5	
Pill	1.716	36.2	49.5	15.4	24.6	3.2	4.5	6.1	8.8	17.0	23.1	
IUD	429	15.2	23.4	0.8	5.0	3.7	4.0	3.6	7.1	7.9	9.6	
Injectable	612	36.6	56.5	9.5	19.1	11	2.5	118	20.8	19.6	31.1	
Condom	210	72.5	81.5	50.8	61.6	8.5	18.4		0.0	38.7	40.7	
Traditional methods8	183	42.0	56.5	21.5	33.6	19.2	25.3	0.4	1.3	8.1	11.2	

Table 2. Cumulative 12- and 24-month probabilities of discontinuation per 100 episodes of method use, by country and method, according to cause

*Vaginal methods, injectable and condom. †Injectable and condom. ‡Vaginal methods and injectable. §Periodic abstinence and withdrawal. Note: Episodes are weighted by sample weight where necessary.