

Table 4. Health professionals' estimates of the percentage of women having an abortion who have a complication and the percentage of those with a complication who are hospitalized, by poverty status and residence, according to geographic area

Geographic area	% with complication					% with complications hospitalized				
	All	Nonpoor urban	Nonpoor rural	Poor urban	Poor rural	All	Nonpoor urban	Nonpoor rural	Poor urban	Poor rural
Regional total	34	19	31	31	39	47	64	57	53	40
Subregion										
South Central	34	17	32	31	40	46	66	57	52	37
Southeast	32	22	30	29	37	51	58	58	55	46
Country										
Afghanistan/ Iran/Pakistan	29	19	27	33	30	34	50	44	44	24
Bangladesh	40	21	34	36	43	54	63	58	52	53
India	34	14	31	28	41	47	70	59	53	37
Indonesia	39	27	36	31	44	34	53	52	43	25
Myanmar/Laos/ Cambodia	46	28	39	45	49	58	54	52	59	59
Nepal	42	26	36	39	45	46	53	54	59	41
Philippines	42	32	41	47	51	65	69	69	63	59
Sri Lanka/Malaysia	25	7	24	25	33	88	91	85	94	84
Thailand	34	12	27	28	38	66	70	75	69	64
Vietnam	11	12	8	6	12	40	33	38	37	43