

**Table 5. Percentage (and standard error) of sexually active high school adolescents aged 15–17 who reported using oral contraceptives at last intercourse, who reported using condoms at last intercourse or who had two or more sexual partners in the past three months, by survey, according to gender**

Survey, year and comparison	Used oral contraceptives		Used condoms		Had ≥2 partners	
	Females	Males	Females	Males	Females	Males
<b>YRBS</b>						
1991	23.6 (3.0)	13.9 (1.2)	39.7 (3.0)	57.4 (2.1)	18.0 (1.3)	39.1 (2.1)
1993	19.4 (1.1)	11.7 (1.1)	47.8 (1.4)	61.6 (2.1)	19.2 (1.3)	39.6 (1.9)
1995	18.3 (1.7)	11.7 (1.9)	49.2 (2.8)	61.6 (2.8)	16.1 (1.8)	37.5 (2.8)
1997	17.6 (1.8)	11.3 (1.6)	51.9 (1.7)	63.8 (1.4)	19.2 (1.3)	34.0 (1.9)
Trend over time	ns	ns	p<.001	p<.05	ns	p<.05
<b>NSFG/NSAM</b>						
1988	33.8 (3.7)	27.6 (3.4)	38.4 (4.3)	61.5 (4.3)	8.7 (2.6)	19.0 (3.1)
1995	20.7 (3.2)	24.7 (4.4)	41.5 (3.6)	72.1 (3.2)	15.3 (2.8)	20.5 (3.4)
Trend over time	p<.01	ns	ns	p<.05	ns	ns
<b>Add Health</b>						
1995	25.6 (1.8)	20.1 (1.7)	53.1 (2.0)	64.2 (2.3)	u	u
<b>Differences between surveys in 1995</b>						
YRBS vs. NSFG/NSAM	ns	p<.01	ns	p<.05	ns	p<.001
NSFG/NSAM vs. Add Health	ns	ns	p<.01	p<.05	na	na
YRBS vs. Add Health	p<.01	p<.01	ns	ns	na	na

*Notes:* The denominator for analyses includes all high school–attending adolescents aged 15–17 who reported having had sexual intercourse in the past three months and who completed the survey. Estimates for racial and ethnic subgroups are not presented, as multiple cells contained less than 100 respondents. Comparable data on number of partners are not available from Add Health. ns=non-significant. na=not applicable. u=unavailable.