

**TABLE 5. Percentages (and 95% confidence intervals) of women seeking postabortion care who received selected types of care, and percentage distribution of these women by type of provider—all according to type of facility**

Measure	All (N=57,964)	Public hospital (N=20,198)	Public health center (N=23,317)	Private/NGO facility (N=14,449)
<b>Referred elsewhere***</b>	13 (9–16)	1 (1–2)	27 (19–35)	4 (1–7)
<b>Evacuation performed</b>				
Any***	79 (75–83)	93 (91–95)	59 (50–68)	95 (92–98)
Sharp curettage†,***	38	54	25	25
MVA/EVA†,***	56	37	69	72
Other†,‡	6	9	6	3
<b>Antibiotics provided</b>	83 (79–87)	85 (79–91)	79 (73–86)	86 (79–93)
<b>Transfusion administered*</b>	2 (1–3)	4 (2–5)	1 (0–1)	1 (0–2)
<b>Intravenous fluids given***</b>	44 (39–49)	66 (61–72)	35 (27–43)	19 (11–26)
<b>Pain medication given***</b>	54 (48–59)	60 (51–68)	40 (30–50)	71 (58–85)
<b>Hospitalization required***</b>	23 (19–26)	43 (38–49)	9 (5–13)	11 (6–16)
<b>Evacuation providers§,***</b>				
Physician	43 (37–49)	68 (70–76)	3 (1–6)	81 (72–90)
Midlevel provider	35 (30–41)	25 (18–32)	55 (45–64)	14 (6–22)
Missing	21 (17–26)	7 (5–10)	42 (33–51)	5 (2–8)

\*Percentages differ among facility types at  $p < .05$ . \*\*\*Percentages differ among facility types at  $p < .001$ .  
 †Because of the large number of strata used in the analysis, the standard error estimates for the “other” subcategory are unstable; thus, only point estimates are shown for the three subcategories. ‡Includes pitocin, medical methods and manual removal of retained products. §Specialists, general practitioners, residents and interns are considered physicians; nurses, midwives and health officers are midlevel providers. Notes: All percentages are weighted and unless otherwise noted exclude respondents with missing data. Hospitalization refers to a stay of at least 24 hours. NGO=nongovernmental organization. MVA/EVA=manual vacuum aspiration or electric vacuum aspiration.