

**TABLE 4. Percentage of women aged 15–49 currently in union using a contraceptive method, by method category and year, and percentage with unmet need, by year—all according to region, Colombia**

Region	% using a method								% with unmet need‡			
	Modern*				Traditional†							
	1990	2000	2005	2010	1990	2000	2005	2010	1990	2000	2005	2010
<b>All</b>	<b>54.6</b>	<b>63.3</b>	<b>67.5</b>	<b>72.8</b>	<b>11.5</b>	<b>13.6</b>	<b>10.7</b>	<b>6.3</b>	<b>11.1</b>	<b>6.2</b>	<b>5.8</b>	<b>6.9</b>
Atlántica	48.4	57.3	59.0	65.0	5.0	13.5	11.4	6.5	18.3	9.0	9.6	10.2
Bogotá	61.8	68.7	74.1	75.7	13.0	11.9	8.4	5.1	7.9	4.5	3.6	5.4
Central	53.1	62.8	69.4	75.3	12.4	13.9	10.4	6.4	9.6	5.5	4.9	5.4
Oriental	57.0	66.4	69.6	75.2	16.7	16.1	11.8	6.3	7.5	3.7	4.4	6.0
Pacífica	54.8	64.2	66.9	73.5	10.9	12.1	11.3	6.9	11.8	7.6	5.5	7.7

\*Includes condom, implant, injectable, IUD, pill, spermicide and female or male sterilization. †Includes periodic abstinence, withdrawal, lactational amenorrhea and folk methods. ‡Percentage who are fecund, do not want a child in the next two years (or ever) and are not using any method of contraception. Sources: Special tabulations of data from Demographic and Health Surveys (references 19–22).