

TABLE 3. Percentage distribution of sexually active female high school students, by contraceptive method used at last sexual intercourse, and pregnancy risk score, both according to race or ethnicity, 1991 and 2003

Method	1991			2003		
	White (N=927)	Black (N=811)	Hispanic (N=443)	White (N=979)	Black (N=744)	Hispanic (N=678)
Pill only	24.6 (3.0)	18.3 (2.1)	16.5 (2.6)	18.4 (1.6)	8.5 (1.7)	8.3 (1.9)
Condom only	35.0 (2.8)	35.3 (2.6)	26.2 (2.9)	47.5 (1.7)	56.1 (2.6)	47.3 (4.2)
Pill and condom	3.0 (0.8)	4.2 (1.1)	0.2 (0.2)	7.7 (1.1)	3.0 (0.9)	3.7 (1.5)
Withdrawal	21.4 (2.4)	13.6 (2.0)	19.2 (3.1)	11.7 (1.6)	7.2 (1.3)	12.2 (1.8)
Injectable	u	u	u	2.9 (0.7)	5.2 (1.3)	2.3 (0.9)
Injectable and condom	u	u	u	1.3 (0.3)	4.0 (1.2)	1.1 (0.6)
Other method	2.1 (0.6)	2.5 (0.9)	4.8 (2.6)	1.6 (0.6)	2.2 (0.7)	2.7 (0.7)
Other and condom	0.1 (0.1)	0.1 (0.1)	0.5 (0.3)	0.3 (0.2)	0.6 (0.4)	0.1 (0.1)
No method	13.4 (1.4)	25.6 (1.9)	28.8 (3.3)	8.5 (1.1)	11.6 (2.0)	21.4 (3.5)
Not sure	0.5 (0.3)	0.4 (0.3)	3.8 (2.0)	0.1 (0.1)	1.8 (0.6)	0.8 (0.4)
Total	100.0	100.0	100.0	100.0	100.0	100.0
<i>Pregnancy risk score</i>	22.8 (1.0)	37.9 (1.3)	39.3 (2.8)	17.2 (0.9)***	29.0 (1.4)***	31.8 (2.5)*

*Trend in pregnancy risk from 1991 to 2003 significant at p<.05. ***Trend in pregnancy risk from 1991 to 2003 significant at p<.001. Notes: Figures in parentheses are standard errors. The pregnancy risk score is based on method-specific contraceptive failure rates and the proportion using each method. u=unavailable. Percentages may not total 100.0 because of rounding. White and black include non-Hispanics only.