

**TABLE 1. Among married and sexually active unmarried women aged 15–29, percentage currently using a modern contraceptive method; percentage distribution of those currently using a contraceptive, by method; and percentage with unmet need for spacing or limiting—Urban Reproductive Health Initiative, Senegal, 2011**

Marital status and age	% using modern method	% distribution of users							% unmet need		
		Pill	Injectable	Implant	IUD	Condom	Other	Total	Spacing	Limiting	Total
<b>Married (N=2,340)</b>											
All	19.7 (18.0–21.6)	32.7 (28.1–37.6)	42.5 (37.6–47.5)	6.9 (4.7–10.0)	2.3 (1.3–4.1)	15.2 (11.7–19.3)	0.5 (0.1–1.4)	100.0	18.9 (17.2–20.8)	0.3 (0.1–0.5)	19.2 (17.5–21.1)
15–19	7.6 (5.1–11.1)	30.5 (14.3–53.5)	40.7 (23.6–60.4)	3.7 (0.7–17.1)	0.0 na	25.1 (12.0–45.3)	0.0 na	100.0	15.3 (11.3–20.4)	0.0 na	15.3 (11.3–20.4)
20–24	21.6 (18.7–24.7)	28.6 (22.1–36.1)	44.6 (37.1–52.4)	3.8 (1.9–7.4)	2.8 (1.2–6.3)	20.3 (14.2–28.1)	0.0 na	100.0	20.3 (17.6–23.3)	0.2 (0.1–0.5)	20.5 (17.9–23.5)
25–29	21.7 (19.1–24.5)	36.3 (29.9–43.3)	40.9 (34.2–48.0)	9.7 (6.1–15.2)	2.2 (1.0–4.8)	10.0 (6.7–14.6)	0.9 (0.3–2.7)	100.0	18.8 (16.3–21.7)	0.4 (0.2–0.8)	19.2 (16.6–22.0)
<b>Unmarried (N=237)</b>											
All	27.1 (20.7–34.7)	14.1 (6.6–27.4)	20.8 (11.9–33.8)	7.3 (1.6–27.3)	0.6 (0.1–3.9)	55.8 (40.3–70.2)	1.5 (0.2–9.8)	100.0	10.6 (6.7–16.5)	0.0 na	10.6 (6.7–16.5)

Notes: Figures in parentheses are 95% confidence intervals. na=not applicable.

**TABLE 2. Percentage distribution (and 95% confidence intervals) of currently married and sexually active unmarried young women using a modern contraceptive method, by source of most recent method, according to method**

Source	All (N=536)	Pill (N=166)	Injectable (N=239)	Implant (N=36)	Condom (N=77)
<b>Public</b>	<b>66.5</b>	<b>66.1 (57.3–73.9)</b>	<b>83.8 (76.0–89.4)</b>	<b>95.4 (83.3–98.9)</b>	<b>5.2 (1.8–13.7)</b>
Hospital	2.1	2.3 (0.8–6.2)	1.5 (0.6–3.8)	10.4 (4.4–22.5)	0.0
Health center	17.2	19.3 (13.1–27.7)	13.3 (9.2–18.9)	60.1 (41.1–76.6)	1.2 (0.2–8.1)
Health post	42.8	39.3 (30.9–48.4)	63.7 (55.9–70.9)	24.1 (11.8–43.0)	3.7 (1.0–12.5)
Other public†	4.4	5.2 (2.3–11.2)	5.3 (2.9–9.6)	0.8 (0.1–5.8)	0.2 (0.0–1.7)
<b>Private</b>	<b>26.1</b>	<b>29.9 (22.5–38.5)</b>	<b>14.6 (9.2–22.5)</b>	<b>0.0</b>	<b>61.1 (46.7–73.8)</b>
Hospital/clinic	8.2	9.1 (4.9–16.2)	11.3 (6.4–19.3)	0.0	1.0 (0.1–6.7)
Other private	17.9	20.8 (14.8–28.5)	3.3 (1.5–7.2)	0.0	60.1 (45.8–72.9)
<b>NGO/other‡</b>	<b>7.4</b>	<b>4.0 (1.6–9.7)</b>	<b>1.6 (0.6–4.1)</b>	<b>4.6 (1.1–16.7)</b>	<b>33.7 (21.5–48.5)</b>
Total	100.0	100.0	100.0	100.0	100.0

†Includes dispensaries and community health centers. ‡Includes workplace clinics, youth centers, voluntary counseling and testing centers, shops, markets and peer educators. Notes: NGO=nongovernmental organization. Data for the IUD and other methods are not presented because of small sample sizes.

**TABLE 3. Percentage of family planning providers who apply a minimum age or marital status restriction to contraceptive provision, by method; and median minimum age restriction (and interquartile range), by method—all according to facility type**

Barrier/method	Public					Private
	All (N=516)	Hospitals (N=32)	Health centers (N=81)	Health posts (N=364)	Other† (N=39)	All (N=121)
<b>Minimum age</b>						
Pill	57.0	59.3	47.3	46.0	46.9	48.8
Injectable	43.6	51.9	42.5	40.1	36.7	41.4
Implant	45.2	52.0	44.9	32.7	37.5	38.1
Condoms	24.5	29.6	25.0	16.0	25.7	19.9
EC	24.2	26.9	22.2	18.1	34.8	21.1
<b>Median minimum age</b>						
Pill	17 (3)	18 (4)	17 (3)	17 (3)	15 (2)	18 (3)
Injectable	18 (5)	19 (2)	18 (9)	18 (5)	18 (7)	18 (4)
Implant	18 (7)	20 (7)	20 (7)	18 (9)	18 (2)	18 (4)
Condoms	18 (3)	18 (5)	17 (3)	18 (3)	18 (1)	18 (2)
EC	18 (3)	18 (5)	17 (5)	17 (3)	18 (2)	18 (3)
<b>Marital status</b>						
Pill	12.2	12.5	18.5	9.6	23.1	20.7
Injectable	13.6	12.5	18.5	12.1	17.9	28.1
Implant	13.6	12.5	25.9	11.0	12.8	29.8
Condoms	8.3	6.3	11.1	7.4	12.8	11.6
EC	8.7	6.3	13.6	7.7	10.3	21.5

†Includes dispensaries and community health centers. Note: EC=emergency contraception.

**TABLE 4. Percentage of family planning providers who apply a minimum age restriction to contraceptive provision, by providers' gender, age and type, according to health care sector and method**

Characteristics	Public					Private				
	No. of providers	Pill	Injectable	Condom	Any of the 3	No. of providers	Pill	Injectable	Condom	Any of the 3
<b>Sex</b>										
Male	86	65.2	54.1**	32.0	58.1*	21	66.7	63.2	30.0	66.5
Female	430	44.0	38.9**	16.4	44.7*	100	54.9	39.0	23.2	51.0
<b>Age</b>										
<30	66	47.5	42.6	18.0	48.5	22	55.0	45.0	31.6	54.5
30–39	153	44.2	41.2	21.0	48.4	40	52.9	38.2	23.5	52.5
≥40	297	48.5	40.5	18.0	45.8	59	60.9	46.8	22.4	54.2
<b>Provider type</b>										
Doctor	13	33.3**	45.5	33.3	38.5**	19	52.9	44.4	16.7	57.9
Nurse	131	62.0**	51.9	31.0	57.3**	25	70.6	52.9	40.0	60.0
Midwife/other	372	42.8**	37.5	14.2	43.5**	77	54.5	40.9	21.9	50.6

\*Differences across subgroups significant at  $p < .05$ . \*\*Differences across subgroups significant at  $p < .01$ . Notes: Differences assessed using chi-square tests. Few providers of the pill, the injectable and condoms in the public sector were male.

**TABLE 5. Percentage of family planning providers who apply a marital status restriction to contraceptive provision, by providers' gender, age and type, according to health care sector and method**

Characteristics	Public					Private				
	No. of providers	Pill	Injectable	Condom	Any of the 3	No. of providers	Pill	Injectable	Condom	Any of the 3
<b>Sex</b>										
Male	86	9.3	9.3	8.1	12.8	21	19.0	14.3	14.3	19.0
Female	430	12.8	14.4	8.4	18.1	100	21.0	31.0	11.0	36.0
<b>Age</b>										
<30	66	4.5*	6.1	6.1	7.6	22	31.8	40.9	9.1	45.5
30-39	153	9.2*	11.8	5.9	17.0	40	20.0	35.0	10.0	37.5
≥40	297	15.5*	16.2	10.1	19.5	59	16.9	18.6	13.6	25.4
<b>Provider type</b>										
Doctor	13	7.7	7.7	7.7	7.7	19	21.1	15.8	15.7	21.1
Nurse	131	14.5	13.9	9.2	17.5	25	28.0	28.0	16.0	32.0
Midwife/other	372	1.6	13.7	8.1	17.7	77	18.2	31.2	9.1	36.4

\*Differences across age-groups significant at  $p < .05$ . Notes: Differences assessed using chi-square tests. Few providers of the pill, the injectable and condoms in the public sector were male.