

TABLE 1. Percentage distribution of sexually experienced female adolescent clinic attendees, by selected characteristics, Greater Boston, 2006

Characteristic	% (N=356)
Age	
14–15	14
16–17	47
18–20	39
Race or ethnicity	
White	40
Black	20
Hispanic	34
Other	6
Education	
Enrolled in high school	62
GED program/high school dropout	7
≥high school	32
Living situation	
With parents or family	83
With partner	7
Other*	10
Experienced partner abuse	
Yes	45
No	55
Total	100

*With friends, between homes, with child protection services or in foster care.

Note: Percentages may not total 100 because of rounding.

TABLE 2. Percentage of women reporting standard sexual risk behaviors and coercive or deceptive sexual risk factors, by lifetime history of intimate partner violence, and odds ratios (and 95% confidence intervals) from logistic regression analyses examining associations between sexual risk behaviors and history of violence

Sexual risk behavior or factor	All	Those reporting violence	Those reporting no violence	Odds ratio*
Standard				
>1 vaginal sex partnert	21	27	15	2.2 (1.3–3.7)
Unprotected vaginal sext	64	62	65	1.0 (0.6–1.5)
Anal sex	27	33	23	1.7 (1.1–2.7)
Unprotected anal sext†	11	15	8	2.2 (1.1–4.3)
Coercive/deceptive				
Male partner infidelity	42	58	29	3.4 (2.2–5.4)
Fear of requesting condom use	12	18	7	2.9 (1.4–6.0)
Experienced negative consequence of condom request	12	21	5	5.3 (2.4–11.6)
Coerced into not using a condom	19	31	9	4.9 (2.6–8.9)
Forced anal sex	1	3	<1	‡

*Adjusted for age, race or ethnicity, and recruitment site. †In the past three months. ‡Because of small cell size, logistic regression was not performed; however, Fisher's exact test showed significant difference between those reporting violence and those reporting no violence ($p < .05$).