

**TABLE 1. Selected characteristics of adolescents aged 16 or older at Wave 2, National Longitudinal Study of Adolescent Health**

Characteristic	% or mean (N=5,798)
<b>Individual/family†</b>	
Male	51.1
Mean age	16.2 (1.1)
Race/ethnicity	
White	65.0
Black	16.7
Hispanic	12.1
Other	6.2
Lived with two biological/adoptive parents	53.6
Mean parent-teenager closeness (range, 0–4)	3.2 (0.8)
Ever used substances	70.4
Mean PVT score (range, 13–146)	101.6 (14.8)
Had high educational aspirations	66.0
Education at Wave 3	
<high school	10.1
High school	37.0
≥some college	52.8
Parent's education	
<high school	13.5
High school	31.3
≥some college	55.3
<b>Sexual risk factors‡</b>	
Mean risk index (range, 0–8)	2.1 (2.4)
No. of risk factors	
0	42.5
1–2	18.2
3–4	22.3
5–8	17.0
Inconsistent contraceptive use	34.5
First sex before age 16	26.2
Ever had older partner	18.1
Ever did not discuss contraception before first sex with a romantic partner§	29.0
Multiple partners by Wave 2	42.0
Ever had nonmonogamous partner	17.6
Ever had nonromantic partner	20.3
Ever had one-night stand	21.5
Had not had sex by Wave 2	38.5
<b>Dependent variables</b>	
No. of partners in last year††	
0	8.4
1	59.6
≥2	32.0
Had STD in last year††	16.9
Had a birth/fathered a child at age 20–24‡‡	
No	75.2
Intended	9.6
Unintended	15.2

†Measured at Wave 1 (1995) unless noted otherwise. ‡Measured at Waves 1 and 2 (1996) unless noted otherwise. §Because this measure applied only to romantic relationships, 5% of respondents did not answer it. ††Measured at Wave 3 (2002). ‡‡Measured at Wave 4 (2008). Notes: Figures are weighted percentages unless noted otherwise. For means, numbers in parentheses are standard deviations. PVT=Peabody Picture Vocabulary Test.

**TABLE 2. Relative risk ratios and odds ratios from multivariate logistic regression analyses assessing associations between selected characteristics and reproductive health and behavior outcomes in young adulthood**

Characteristic	No. of partners in last year† (N=5,743)			STD diagnosis in last year† (N=4,898)	Birth at age 20–24† (N=5,036)		
	1 (vs. 0)	≥2 (vs. 0)	≥2 (vs. 1)		Intended (vs. none)	Unintended (vs. none)	Intended (vs. unintended)
<b>MODEL 1</b>							
<b>Cumulative risk index</b>	1.2**	1.3***	1.2***	1.1***	1.1**	1.1***	1.0
<b>MODEL 2</b>							
<b>Individual sexual risk factors</b>							
Inconsistent contraceptive use	1.2	1.6*	1.3*	1.2	1.2	1.5**	1.3
First sex before age 16	0.9	0.8	0.9	1.3	0.6*	0.8	1.3
Ever had older partner	1.0	1.1	1.1	1.0	1.1	1.0	0.9
Ever did not discuss contraception before first sex with a romantic partner	1.3	1.5	1.2	0.9	1.2	1.3*	1.1
Multiple partners by Wave 2	1.0	1.3	1.3*	1.0	1.4	1.3	0.9
Ever had nonmonogamous partner	1.6*	2.0**	1.2	1.1	1.2	0.9	0.8
Ever had nonromantic partner	1.4	1.9*	1.4**	1.3	1.2	1.1	0.9
Ever had one-night stand	1.0	0.9	0.9	1.0	1.1	1.0	0.9
<b>Individual/family</b>							
Male	0.6***	1.2	2.0***	0.5***	0.4***	0.5***	1.2
Age at Wave 1	1.0	0.8**	0.8***	1.0	0.8*	0.9	1.1
Race/ethnicity							
White (ref)	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Black	1.4	3.0***	2.2***	3.8***	1.0	1.4*	1.4
Hispanic	1.6*	1.5	1.0	1.1	1.1	0.8	0.8
Other	1.1	1.1	1.0	1.5*	0.9	0.7	0.8
Lived with two parents	0.8	1.0	1.2	0.9	0.7**	0.6***	0.9
Parent-teenager closeness	1.1	1.1	0.9	0.9	1.0	0.9	0.9
Ever used substances	1.3	1.7**	1.3*	1.0	0.9	0.9	1.0
PVT score	1.0	1.0	1.0	1.0	1.0	1.0	1.0
High educational aspirations	1.1	1.0	0.9	0.9	0.8	0.8	1.0
Education at Wave 3							
<high school	1.2	1.1	1.0	1.0	1.2	1.0	0.8
High school (ref)	1.0	1.0	1.0	1.0	1.0	1.0	1.0
≥some college	0.9	1.2	1.4**	1.0	0.4***	0.4***	1.0
Parent's education							
<high school	1.0	1.0	1.1	1.1	0.9	1.0	1.2
High school (ref)	1.0	1.0	1.0	1.0	1.0	1.0	1.0
≥some college	1.1	1.2	1.0	1.1	0.7**	0.9	1.3

\*p<.05. \*\*p<.01. \*\*\*p<.001. †Relative risk ratios. ‡Odds ratios. Notes: Model 1 included all of the same controls as Model 2; results for controls were virtually the same in both models. ref=reference category. PVT=Peabody Picture Vocabulary Test.