

**TABLE 1. Percentage distribution of physicians participating in a study of contraceptive recommendations for standardized patients, by selected characteristics, 2007–2008**

Characteristic	% (N=468)
<b>Sex</b>	
Male	55
Female	45
<b>Race/ethnicity</b>	
White	77
Black	8
Hispanic	4
Asian	10
Other	2
<b>Age</b>	
≤35	18
36–45	33
46–55	31
≥56	18
<b>Specialty</b>	
Obstetrics-gynecology	62
Family medicine	38
<b>Frequency of prescribing contraceptives</b>	
Frequently	83
Occasionally	17
<b>Board-certified</b>	
Yes	92
No	8
<b>Practice type</b>	
Academic	25
Private	54
HMO	8
Other‡	13
<b>Region</b>	
Northeast	18
Midwest	32
South	31
West	19
<b>Inserts IUDs</b>	
Yes	79
No	21
Total	100

‡Family planning clinic or community health center. Note: Percentages may not total 100 because of rounding.

**TABLE 2. Percentage of physicians recommending specific contraceptive methods, by selected characteristics**

Characteristic	Pill	Ring	Levonorgestrel IUD	Patch	Injectable	Copper IUD
<b>All</b>	<b>89</b>	<b>80</b>	<b>64</b>	<b>56</b>	<b>49</b>	<b>45</b>
<b>Sex</b>						
Male	87	73***	62	54	50	44
Female	91	89	66	58	49	46
<b>Race/ethnicity</b>						
White	89	81	66*	56	48	44
Black	90	82	71	45	58	55
Hispanic	88	59	71	65	59	59
Asian	89	78	49	60	49	40
Other	70	60	30	50	60	30
<b>Age</b>						
≤35	91	89**	79**	58	42	57
36–45	91	84	67	56	52	44
46–55	85	77	56	57	54	41
≥56	88	67	58	51	44	41
<b>Specialty</b>						
Obstetrics-gynecology	89	83*	70***	50**	43***	49*
Family medicine	88	74	54	66	60	38
<b>Frequency of prescribing contraceptives</b>						
Frequently	90*	84***	68***	55	47*	47*
Occasionally	81	58	47	61	60	33
<b>Board-certified</b>						
Yes	90**	79	64	55	49	45
No	76	87	70	65	49	49
<b>Practice type</b>						
Academic	82*	79	78**	52	47	57**
Private	93	83	61	57	47	38
HMO	83	78	64	58	50	53
Other‡	87	70	51	56	60	44
<b>Region</b>						
Northeast	88	85	63	50	50	44
Midwest	89	79	68	51	50	48*
South	90	76	59	61	42	35
West	87	81	67	60	58	56
<b>Inserts IUDs</b>						
Yes	88	82*	70***	54	46*	49***
No	92	71	41	63	60	29

\*p<.05. \*\*p<.01. \*\*\*p<.001. ‡Family planning clinic or community health center. Note: All p values represent overall group differences.

**TABLE 3. Odds ratios (and 95% confidence intervals) from logistic regression analyses examining associations between physicians' characteristics and their likelihood of recommending specific contraceptive methods**

Characteristic	Pill	Ring	Levonorgestrel IUD	Patch	Injectable	Copper IUD
<b>Sex</b>						
Male (ref)	1.0	1.0	1.0	1.0	1.0	1.0
Female	1.0 (0.6–1.7)	2.1 (1.3–3.6)**	0.9 (0.5–1.4)	1.0 (0.6–1.5)	0.7 (0.4–1.1)†	0.6 (0.4–1.0)†
<b>Race/ethnicity</b>						
White (ref)	1.0	1.0	1.0	1.0	1.0	1.0
Black	0.6 (0.2–1.4)	0.3 (0.1–0.6)**	0.9 (0.4–2.0)	0.6 (0.3–1.4)	1.2 (0.5–2.6)	1.5 (0.7–3.3)
Hispanic	0.9 (0.2–3.0)	0.3 (0.1–0.8)*	2.1 (0.6–6.5)	1.3 (0.5–3.8)	0.9 (0.3–2.9)	3.0 (1.0–8.7)*
Asian	0.5 (0.2–1.2)	0.4 (0.2–0.9)*	0.5 (0.3–1.1)†	1.2 (0.6–2.4)	0.8 (0.4–1.7)	1.0 (0.5–2.2)
Other	0.3 (0.1–1.2)†	0.6 (0.1–2.6)	0.3 (0.1–1.4)	1.4 (0.3–5.3)	2.2 (0.6–8.4)	0.2 (0.03–2.1)
<b>Age</b>						
≤35 (ref)	1.0	1.0	1.0	1.0	1.0	1.0
36–45	0.9 (0.4–2.0)	0.5 (0.2–1.2)	0.4 (0.2–0.7)**	1.1 (0.6–2.1)	1.9 (0.9–3.9)†	0.5 (0.2–0.9)*
46–55	0.7 (0.3–1.6)	0.5 (0.2–1.1)†	0.3 (0.1–0.6)**	1.1 (0.6–2.2)	1.8 (0.9–3.9)	0.3 (0.2–0.7)**
≥56	1.0 (0.4–2.5)	0.3 (0.1–0.6)**	0.2 (0.2–0.5)**	1.1 (0.5–2.3)	1.9 (0.8–4.4)	0.4 (0.2–0.9)*
<b>Specialty</b>						
Obstetrics-gynecology (ref)	1.0	1.0	1.0	1.0	1.0	1.0
Family medicine	0.8 (0.4–1.5)	0.6 (0.3–1.0)*	0.8 (0.5–1.3)	2.6 (1.6–4.4)***	2.5 (1.5–4.3)**	0.9 (0.5–1.6)
<b>Frequency of prescribing contraceptives</b>						
Frequently (ref)	1.0	1.0	1.0	1.0	1.0	1.0
Occasionally	0.7 (0.4–1.4)	0.6 (0.3–1.0)†	0.6 (0.3–1.1)	1.4 (0.8–2.6)	1.7 (0.9–2.9)†	0.6 (0.3–1.2)
<b>Board-certified</b>						
Yes	2.1 (0.9–5.0)†	0.9 (0.3–2.5)	1.2 (0.5–3.0)	0.6 (0.3–1.5)	0.6 (0.2–1.4)	1.8 (0.8–4.4)
No (ref)	1.0	1.0	1.0	1.0	1.0	1.0
<b>Practice type</b>						
Academic (ref)	1.0	1.0	1.0	1.0	1.0	1.0
Private	2.9 (1.7–5.0)***	2.4 (1.4–4.2)**	0.6 (0.4–1.0)†	1.5 (0.9–2.5)	0.9 (0.6–1.6)	0.5 (0.3–0.9)*
HMO	1.5 (0.6–3.7)	2.1 (0.8–5.4)	0.6 (0.2–1.4)	2.0 (0.8–4.6)	0.4 (0.2–1.1)	0.5 (0.2–1.2)
Other†	2.0 (0.9–4.5)†	1.1 (0.5–2.3)	0.5 (0.2–1.0)†	1.9 (0.9–3.8)†	1.4 (0.7–2.8)	0.8 (0.4–1.6)
<b>Region</b>						
Northeast (ref)	1.0	1.0	1.0	1.0	1.0	1.0
Midwest	1.2 (0.6–2.4)	0.6 (0.3–1.2)	1.0 (0.5–1.8)	1.3 (0.7–2.5)	0.8 (0.4–1.5)	1.3 (0.7–2.5)
South	1.3 (0.6–2.5)	0.8 (0.4–1.5)	0.6 (0.4–1.2)	2.6 (1.4–4.9)**	0.6 (0.3–1.1)	0.7 (0.4–1.4)
West	0.7 (0.3–1.5)	0.6 (0.3–1.4)	0.8 (0.4–1.5)	1.3 (0.7–2.7)	1.2 (0.6–2.4)	1.6 (0.8–3.2)
<b>Inserts IUDs</b>						
Yes	0.4 (0.2–0.8)*	0.8 (0.4–1.4)	2.2 (1.2–4.1)*	1.0 (0.6–1.9)	0.8 (0.4–1.4)	1.7 (0.9–3.4)
No (ref)	1.0	1.0	1.0	1.0	1.0	1.0

\*p<.05. \*\*p<.01. \*\*\*p<.001. †p<.10. ‡Family planning clinic or community health center. Notes: A physician was considered to have recommended a method if his or her score for that method was higher than the physician's mean recommendation score for all six methods. All models control for the standardized patient's race and ethnicity, socioeconomic status and gynecologic history; the model for the levonorgestrel IUD also includes an interaction term between patient's race and ethnicity and socioeconomic status. ref=reference group.