

TABLE 2. Mean number of days during which nonpregnant study participants did not use an effective contraceptive method, by number of months since first birth, according to study group

Months since first birth	All methods		LARC use removed	
	Enhanced care	Usual care	Enhanced care	Usual care
0–3	52.6 (48.3–57.3)**	62.2 (58.2–66.6)	65.0 (56.1–75.3)**	79.2 (70.1–89.3)
4–6	20.5 (15.8–26.6)	28.0 (22.6–34.6)	39.2 (30.9–49.7)*	54.9 (45.2–66.5)
7–9	22.6 (17.5–29.2)	27.1 (21.5–34.1)	43.3 (34.2–54.8)	54.4 (45.2–65.5)
10–12	24.4 (19.0–31.3)	29.1 (23.3–36.2)	47.4 (38.0–59.1)	57.9 (48.5–69.2)
13–15	27.3 (21.3–35.0)	33.8 (27.4–41.5)	53.4 (42.9–66.3)	66.2 (56.0–78.3)
16–18	34.8 (28.1–43.1)	31.4 (25.3–39.1)	64.1 (52.5–78.2)	67.4 (56.5–80.4)
19–21	35.5 (28.6–43.9)	35.5 (28.5–44.1)	62.5 (50.7–77.0)	66.2 (55.7–78.8)
22–24	35.5 (28.7–44.0)	36.9 (29.5–46.2)	62.2 (50.5–76.7)	73.2 (61.7–86.8)

*Significantly different from mean number of days for the usual care group at $p < .05$. **Significantly different from mean number of days for the usual care group at $p < .01$. Notes: Survey intervals were standardized to 90 days. Figures in parentheses are 95% confidence intervals. LARC=long-acting reversible contraceptive.