

Table 2. Respondents' estimates of the percentage distribution of women having abortions, by type of provider, according to poverty status, residence and health zone

Poverty status, residence and health zone	N	Physician	Nurse/midwife	Traditional practitioner*	Chemist	Self-induced	Total
Total	61	32	21	16	23	8	100
NONPOOR							
Urban	61	67	14	4	10	5	100
Northeast	17	59	17	6	11	7	100
Northwest	19	66	15	2	11	5	100
Southeast	11	69	11	5	13	3	100
Southwest	14	78	12	2	5	3	100
Rural	51	39	22	14	18	7	100
Northeast	16	28	24	15	21	12	100
Northwest	14	40	22	19	13	7	100
Southeast	8	41	22	12	21	5	100
Southwest	13	54	17	10	15	5	100
POOR							
Urban	61	27	23	16	26	9	100
Northeast	17	33	23	12	23	9	100
Northwest	19	24	22	16	27	12	100
Southeast	11	21	18	20	35	7	100
Southwest	14	29	30	16	18	8	100
Rural	53	14	20	32	24	11	100
Northeast	17	14	17	28	28	14	100
Northwest	15	12	17	37	21	13	100
Southeast	8	8	18	41	26	7	100
Southwest	13	22	31	20	18	10	100

*Includes birth attendants, herbalists, healers and other lay practitioners.