

**Table 3. Percentage of married Mayan women of reproductive age who knew of, had ever used and were currently using a contraceptive, by type of method, according to survey date**

Method	Know		Ever used		Currently using	
	1992	1996	1992	1996	1992	1996
<b>Any method</b>	<b>43.1</b>	<b>95.9***</b>	<b>7.3</b>	<b>24.9***</b>	<b>5.2</b>	<b>18.0***</b>
<b>Modern</b>	<b>41.5</b>	<b>94.5***</b>	<b>5.0</b>	<b>18.5***</b>	<b>3.7</b>	<b>13.5***</b>
Pill	30.2	84.1***	2.6	7.5***	1.1	2.7
Tubal ligation	29.6	86.7***	0.2	4.2***	1.8	5.4
Injectable	16.0	72.4***	1.9	5.5***	0.0	2.5
Vasectomy	9.7	56.6***	0.5	0.4	0.5	0.4
Condom	5.3	33.2***	0.2	4.4***	0.2	0.5
IUD	4.8	35.1***	0.1	1.4**	0.1	0.1
Vaginal tablets	4.1	16.5***	0.1	0.5	0.0	1.8
<b>Traditional</b>	<b>9.5</b>	<b>51.9***</b>	<b>2.7</b>	<b>14.1***</b>	<b>1.5</b>	<b>4.5***</b>
Rhythm	5.2	34.3***	1.8	8.4***	1.1	4.2
Lactational amenorrhea	u	19.7	u	5.9	u	u
Natural teas, herbs	3.2	19.3***	0.5	0.9	0.4	0.3
Withdrawal	1.9	6.7***	0.0	2.4***	0.0	0.0

\*\*Difference by year is statistically significant at  $p \leq .01$ . \*\*\*Difference by year is statistically significant at  $p \leq .001$ .