

Table 2. Percentage of all women in the extended postpartum period who intend to use contraceptives in the next year, by number of months since last birth

Region and country	0–12	0–3	3–6	6–9	9–12	12
Total	38.5	53.6	37.8	33.2	30.9	16.9
Sub-Saharan Africa	40.6	49.5	40.6	38.6	34.2	23.3
Benin	37.1	44.4	39.9	36.5	27.7	23.1
Central African Rep.	48.6	57.9	53.0	42.2	41.3	23.3
Comoros	42.1	50.0	46.3	47.6	26.5	20.2
Côte d'Ivoire	30.7	31.8	27.5	30.4	33.1	14.1
Ghana	40.2	47.2	37.4	42.2	33.1	22.2
Kenya	50.8	66.7	48.8	46.1	45.0	24.1
Mali	42.2	41.4	44.3	39.7	43.0	25.6
Mozambique	34.7	34.7	34.9	36.8	32.7	20.0
Senegal	33.9	42.8	33.8	30.3	28.0	19.5
Uganda	44.1	51.1	43.9	41.9	41.2	30.4
Zambia	51.2	62.6	53.4	47.9	41.4	33.4
Zimbabwe	31.0	63.4	23.5	21.3	17.6	24.2
Middle East	34.8	61.6	30.5	27.0	24.0	10.6
Egypt	37.9	59.0	36.3	28.9	27.9	13.9
Turkey	31.8	64.1	24.6	25.1	20.1	7.4
Asia	38.6	56.2	40.8	29.6	29.8	10.6
Bangladesh	57.6	70.0	61.9	52.3	44.0	13.7
Indonesia	35.3	60.6	35.7	24.5	21.4	8.6
Kazakhstan	38.8	65.5	38.2	20.6	36.1	7.8
Nepal	44.7	57.5	47.8	38.8	35.0	16.5
Philippines	34.9	51.8	36.7	29.4	25.6	9.7
Uzbekistan	20.0	31.6	24.7	12.1	16.7	7.1
Latin America	35.8	56.2	32.6	29.0	28.2	13.1
Bolivia	44.3	57.5	48.4	38.5	35.0	16.8
Brazil	25.3	55.8	18.2	17.8	14.9	6.8
Colombia	24.3	50.6	22.0	17.9	13.3	9.3
Dominican Republic	34.6	62.0	22.4	21.9	31.1	11.4
Guatemala	32.4	44.9	28.4	28.1	28.9	12.0
Haiti	47.1	58.2	45.2	41.9	44.5	23.7
Peru	42.8	64.1	43.7	36.8	29.9	11.7