

Table 5. Percentage of women having a birth who gave birth in a health facility, and percentage who had received tetanus injections prior to delivery, by region and country

Region and country	Delivered in facility	Received tetanus injections
Total	48.7	60.5
Sub-Saharan Africa	45.1	67.4
Benin	64.4	69.3
Central African Rep.	47.0	61.9
Comoros	40.8	50.7
Côte d'Ivoire	42.7	71.0
Ghana	41.9	73.7
Kenya	39.1	87.2
Mali	33.7	48.6
Mozambique	39.1	33.8
Senegal	49.3	82.9
Uganda	33.9	74.1
Zambia	42.7	77.0
Zimbabwe	67.0	78.9
Middle East	51.1	56.6
Egypt	38.0	70.0
Turkey	64.3	43.2
Asia	42.9	64.0
Bangladesh	6.4	74.4
Indonesia	19.2	69.9
Kazakhstan	98.7	u
Nepal	9.4	44.6
Philippines	30.8	67.2
Uzbekistan	92.8	u
Latin America	59.1	65.0
Bolivia	40.7	38.4
Brazil	93.8	56.1
Colombia	80.9	79.0
Dominican Republic	97.5	95.7
Guatemala	34.8	52.6
Haiti	15.1	62.1
Peru	51.2	71.1

Notes: Questions on place of delivery and receipt of tetanus toxoid injections were asked of women who had given birth in a certain number of years preceding the survey; the period ranged from three years to five years, depending on the country. u=unavailable, because it was not asked in the questionnaire.