

**Table 1. Percentage of family planning clients with gonorrhea, chlamydia or trichomoniasis, and bivariate and multivariable odds ratios (with 95% confidence intervals), all by selected characteristics, Kingston, Jamaica, 1995 (N=767)**

Characteristic	Prevalence		Bivariate odds ratio	Multivariable odds ratio
	N	%		
<b>Total</b>	<b>182</b>	<b>23.7</b>	<b>na</b>	<b>na</b>
<b>Age</b>				
≥25	111	20.7	1.0	1.0
<25	71	30.6	1.7 (1.2–2.4)	1.6 (1.1–2.4)
<b>Reported unpleasant vaginal odor</b>				
No	150	22.2	1.0	1.0
Yes	32	34.8	1.9 (1.2–3.0)	1.2 (0.7–2.1)
<b>Reported bleeding/spotting after sex</b>				
No	162	22.7	1.0	1.0
Yes	20	37.0	2.0 (1.1–3.6)	1.5 (0.8–2.8)
<b>Cervical friability</b>				
No	142	21.5	1.0	1.0
Yes	40	37.4	2.2 (1.4–3.4)	1.7 (1.1–2.8)
<b>Urine dipstick</b>				
Neg./trace/+	110	18.5	1.0	1.0
++/+++	71	42.3	3.2 (2.2–4.7)	3.0 (2.0–4.4)
<b>New partner in last 3 months*</b>				
No	158	22.7	1.0	ne
Yes	24	33.8	1.7 (1.0–2.9)	ne
<b>No. of partners in last 3 months*</b>				
1	154	21.9	1.0	ne
≥2	28	43.8	2.8 (1.6–4.7)	ne
<b>No. of partners in last year*</b>				
1	129	20.5	1.0	1.0
≥2	51	38.1	2.4 (1.6–3.5)	2.3 (1.5–3.6)
<b>Has a casual partner*</b>				
No	170	22.9	1.0	ne
Yes	12	48.0	3.1 (1.4–6.9)	ne
<b>Partner may have other partners</b>				
No	33	16.4	1.0	1.0
Yes/do not know	148	26.2	1.8 (1.2–2.8)	1.5 (0.9–2.3)
<b>Partner has urethral discharge</b>				
No	179	23.6	1.0	1.0
Yes	3	37.5	1.9 (0.5–8.2)	1.1 (0.2–5.6)

\*Spearman correlation coefficients ≥0.3 for having a new partner in the last 3 months, number of partners in last 3 months and number of partners in last year, and for number of partners in last 3 months, number of partners in last year and having a casual partner. Notes: na=not applicable; ne=not entered in model.