

**Table 2. Percentage of family planning clients with (gonorrheal or chlamydial) cervical infection, and bivariate and multivariate odds ratios (with 95% confidence intervals), all by selected characteristics**

Characteristic	Prevalence		Bivariate odds ratio	Multivariate odds ratio
	N	%		
<b>Total</b>	<b>108</b>	<b>14.1</b>	<b>na</b>	<b>na</b>
<b>Age</b>				
≥25	63	11.8	1.0	1.0
<25	45	19.4	1.8 (1.2–2.7)	1.8 (1.2–2.8)
<b>Reported unpleasant vaginal odor</b>				
No	90	13.3	1.0	1.0
Yes	18	19.6	1.6 (0.9–2.8)	1.0 (0.6–1.9)
<b>Reported bleeding/spotting after sex</b>				
No	93	13.0	1.0	1.0
Yes	15	27.8	2.6 (1.4–4.8)	2.1 (1.1–4.1)
<b>Cervical friability</b>				
No	80	12.1	1.0	1.0
Yes	28	26.2	2.6 (1.6–4.2)	2.2 (1.3–3.7)
<b>Urine LED</b>				
Neg/trace/+	68	11.4	1.0	1.0
++/+++	39	23.2	2.3 (1.5–3.6)	2.1 (1.3–3.4)
<b>New partner in last 3 months*</b>				
No	97	13.9	1.0	ne
Yes	11	15.5	1.1 (0.6–2.2)	ne
<b>No. of partners in last 3 months*</b>				
1	92	13.1	1.0	ne
≥2	16	25.0	2.2 (1.2–4.1)	ne
<b>No. of partners in last year*</b>				
1	79	12.6	1.0	1.0
≥2	28	20.9	1.8 (1.1–3.0)	1.7 (1.0–2.9)
<b>Has a casual partner*</b>				
No	102	13.7	1.0	ne
Yes	6	24.0	2.0 (0.8–5.1)	ne
<b>Partner may have other partners</b>				
No	21	10.4	1.0	1.0
Yes/d00o not know	86	15.2	1.5 (0.9–2.6)	1.2 (0.7–2.0)
<b>Partner has urethral discharge</b>				
No	105	13.8	1.0	1.0
Yes	3	37.5	3.7 (0.9–15.9)	3.0 (0.6–14.3)

\*Spearman correlation coefficients ≥0.3 for having a new partner in the last 3 months, number of partners in last 3 months and number of partners in last year, and for number of partners in last 3 months, number of partners in last year and having a casual partner. *Notes:* na=not applicable; ne=not entered in model.