

TABLE 4. Percentage of new client visits in which providers addressed specific components of dual-protection counseling

Indicator	Baseline (N=88)	Time 2 (N=76)
HIV/STI risk assessment and risk reduction		
Discussed client's sexual behavior	19.4	34.2*
Discussed partner's sexual behavior	16.4	35.5*
Discussed HIV/AIDS	12.1	42.1*
Discussed client's STI concerns	9.1	21.1*
Indicated to client that she might have an STI	3.6	22.4*
Discussed how to bring up HIV/STI prevention with partner	3.0	23.7*
Dual protection family planning counseling		
Tailored counseling and education to client's personal situation	28.0	67.1*
Compared HIV/STI protective effects of different family planning methods	7.0	42.1*
Showed client how to use the male condom	10.5	34.2*
Showed client how to use the female condom	†	80.3
Discussed how dual protection could be achieved by either one or two family planning methods	4.7	75.0*
Discussed if client's initially preferred method provided protection against HIV/STIs	2.3	49.2*
Discussed with client how she might convince her partner to use a condom	0.0	18.4*
Used dual-protection flip chart	†	47.4
Distributed brochures	3.8	27.6*

*Difference is statistically significant at $p < .05$ by chi-square test. †Female condom and dual-protection flip chart were unavailable at baseline.