

TABLE 1. Means and percentage distributions of respondents, by whether they used dual protection at their most recent sexual encounter, according to selected characteristics

Characteristics	All (N=554)	Dual method use (N=88)	No dual method use (N=466)
MEANS			
Age (mean yrs.)	25.8	25.4	25.9
Schooling (mean yrs.)	9.6	10.7	9.4
PERCENTAGE DISTRIBUTIONS			
Gender			
Female	42.6	48.9	41.4
Male	57.4	51.1	58.6
Previous instruction on condom use from health care worker			
Yes	28.0	40.9	25.5
No	72.0	59.1	74.5
Condom use at last sex			
Yes	50.5	100.0	41.2
No	49.5	0.0	58.8
Other contraceptive used at last sex			
Yes	34.1	100.0	21.7
No	65.9	0.0	78.3
Type of contraceptive used at last sex†			
Injectable	66.7	64.8	68.3
Pill	27.0	27.3	26.7
IUD	1.1	1.1	0.9
Diaphragm	0.5	1.1	0.0
Sterilization	1.6	1.1	2.0
Rhythm	0.5	0.0	0.9
Other/traditional method	2.6	4.5	0.9

†Ns are 189 for all respondents, 88 for dual method use and 101 for no dual method use.