

**Table 1. Percentage distribution or means (and standard deviations) of characteristics of men aged 20–39, by men's STD infection status, 1991 National Survey of Men**

Characteristic	Any STD (N=466)	Gonorrhea (N=298)	No STD (N=2,893)
<b>% DISTRIBUTIONS</b>			
<b>Type of infection</b>			
Viral	31	na	na
Bacterial	69	na	na
<b>Race</b>			
Black	33	49	10
White	67	51	90
<b>Marital status at diagnosis</b>			
Single	89	95	49
Married	11	5	51
<b>Religion</b>			
Any	80	81	89
None	20	19	11
<b>Education</b>			
≤high school	53	64	54
>high school	47	36	46
<b>Used contraceptive at first vaginal sex</b>			
No	74	77	61
Yes	26	23	39
Total	100	100	100
<b>MEANS</b>			
Mos. between diagnosis and interview	9.00 (6.07)	10.93 (5.68)	na
Age at diagnosis	23.16 (5.11)	21.51 (4.35)	na
Age at first sex	14.86 (3.03)	14.91 (3.22)	16.21 (3.66)

*Notes:* In this and subsequent tables, the category "any STD" includes HIV/AIDS, gonorrhea, syphilis, chlamydia, genital herpes and genital warts; na=not applicable.