

**Table 3. Logistic regression coefficients showing the likelihood that men engaged in various types of sexual and health care behavior while infected with an STD, by type of STD and background characteristics of the men**

Characteristic	Any STD			Gonorrhea	
	Had sex	Told partner	Had retest	Had sex	Had retest
Viral STD	1.76**	-0.60	-0.50*	na	na
Mos. between diagnosis and interview	-0.02	0.27**	0.04*	-0.09**	0.03
Age at diagnosis	0.08**	-0.19	-0.05	-0.00	-0.13**
Black	-0.88**	-4.09**	-0.03	-1.30**	-0.47
Single at diagnosis	0.68	-4.74**	-0.93*	-1.37*	-1.55*
Any religion	0.63**	1.84**	1.09**	0.95*	1.68**
>high school education	-0.46*	2.32**	-1.01**	-1.29**	-0.56*
Age at first sex	0.04	-0.21	0.05	-0.10	0.07
Used contraceptive at first sex	-0.22	1.29	-0.08	-0.30	-1.19**
<i>Intercept</i>	<i>-4.71**</i>	<i>11.01*</i>	<i>1.53</i>	<i>1.97</i>	<i>3.50*</i>
<i>-2 log likelihood</i>	<i>399.79</i>	<i>61.01</i>	<i>448.49</i>	<i>195.18</i>	<i>274.43</i>
<i>df</i>	<i>9/436</i>	<i>9/61</i>	<i>9/402</i>	<i>8/277</i>	<i>8/275</i>

\*p≤.10. \*\*p≤.05. Note: Reference categories are not shown.