

**Table 3. Odds ratios (and 95% confidence intervals) for factors associated with condom use, in combination with the pill or alone, among women aged 14–22**

Method and factor	Odds ratio
<b>Pill and condom (N=605)</b>	
Age†	0.85* (0.74–0.98)
Race/ethnicity	
Black	3.65* (2.08–6.43)
Hispanic	1.04 (0.46–2.36)
White	1.00
Risk behavior score‡	0.84 (0.69–1.01)
Age at first intercourse	
≤13 years	0.39* (0.16–0.94)
14–15 years	0.56* (0.33–0.98)
≥16 years	1.00
No. of partners in previous 3 mos.	
0	3.05* (1.31–7.08)
1	1.00
≥2	1.18 (0.52–2.66)
Discussed HIV with adult relatives	2.80* (1.22–6.42)
<b>Condom only (N=1,080)</b>	
Age†	0.85* (0.79–0.93)
Race/ethnicity	
Black	0.87 (0.60–1.26)
Hispanic	0.48* (0.30–0.77)
White	1.00
Risk behavior score‡	0.83* (0.74–0.93)
Age at first intercourse	
≤13	0.27* (0.17–0.45)
14–15	0.65* (0.44–0.94)
≥16	1.00
Seat belt use§	2.51* (1.72–3.70)

\*Significant at  $p < .05$ . †The increase in odds for each one-year increase in age. ‡A seven-point scale with a range of 0–6. Odds ratio reflects the increase in odds for each one-point increase in the score. §Mostly or always vs. sometimes or never.