

**TABLE 2. Percentage of women and men experiencing selected health outcomes, and adjusted relative risk ratios (and 95% confidence intervals) from binomial logistic regression analyses examining the likelihood of those outcomes, by level of intimate partner violence**

Characteristic	%			Relative risk ratios	
	None	Hit	Hit hard	Hit	Hit hard
<b>Women (N=1,665)</b>					
Unhappy with life	11.3	28.5*	37.8*	2.00 (0.83–4.83)	4.11 (1.06–15.93)*,†
Mental distress	41.3	55.8*	68.1*	1.80 (1.13–2.84)*	3.10 (2.25–4.27)*,†
Poor health	31.6	36.2	49.1*,†	1.10 (0.67–1.80)	2.39 (1.10–5.23)*,†
Sexual dysfunction	37.2	41.9	51.8	1.17 (0.68–2.01)	1.95 (1.03–3.67)*
Sexual dissatisfaction					
Physical‡	13.2	16.7	28.1*	1.16 (0.52–2.57)	2.99 (1.46–6.13)*
Emotional‡	8.7	16.1*	28.1*	1.60 (0.62–4.14)	4.40 (2.07–9.33)*
Kissing absent‡	24.0	31.2	38.5	1.30 (0.82–2.06)	3.10 (1.20–8.04)*
Intimacy in sex absent‡	42.2	51.6*	68.0*,†	1.51 (0.86–2.63)	3.46 (2.21–5.39)*,†
Unwanted sexual behavior					
Sex	22.1	30.0*	41.4	1.62 (1.15–2.30)*	2.60 (1.04–6.52)*
Act during sex‡	19.0	31.9*	37.7*	2.18 (1.35–3.51)*	2.60 (1.32–5.13)*
Partner's extramarital sex	8.8	10.8	36.2*,†	1.43 (0.95–2.15)	6.70 (1.85–24.22)*,†
Recent genitourinary symptoms	40.6	50.4*	54.5*	1.54 (1.12–2.12)*	1.67 (1.19–2.33)*
Ever had an STI	1.1	0.3	3.5*,†	0.39 (0.10–1.53)	4.20 (0.89–19.89)†
Positive chlamydia test	3.3	1.0*	1.0*	0.38 (0.15–0.95)*	0.34 (0.13–0.88)*
Lack of AIDS knowledge	54.3	53.4	37.5*,†	0.70 (0.50–0.99)*	0.42 (0.31–0.57)*,†
<b>Men (N=1,658)</b>					
Unhappy with life	10.4	12.4	34.8*	0.93 (0.65–1.35)	5.68 (2.42–13.30)*,†
Mental distress	50.7	54.0	73.6*	1.16 (0.55–2.42)	2.36 (1.12–4.96)*
Poor health	30.5	41.8	55.1*	1.71 (0.67–4.32)	3.62 (1.73–7.57)*
Sexual dysfunction	31.5	36.6	45.0	1.12 (0.62–2.03)	2.53 (0.92–6.92)
Sexual dissatisfaction					
Physical‡	9.0	19.7*	17.8	2.29 (1.01–5.19)*	2.45 (0.60–9.95)
Emotional‡	6.0	21.1*	23.9*	3.87 (2.46–6.08)*	4.90 (2.31–10.39)*
Kissing absent‡	16.6	15.6	8.0	0.73 (0.18–3.01)	0.76 (0.19–3.00)
Intimacy in sex absent‡	62.1	67.8	65.3	1.25 (0.66–2.38)	1.38 (0.62–3.07)
Unwanted sexual behavior					
Sex	6.9	17.1	7.2	2.90 (0.76–11.00)	0.97 (0.26–3.65)
Act during sex‡	10.0	26.4*	11.0	3.19 (1.84–5.52)*	1.11 (0.47–2.61)
Partner's extramarital sex	7.7	18.9	35.0*,†	3.05 (1.06–8.75)*	6.59 (2.52–17.22)*,†
Recent genitourinary symptoms	17.8	16.5	38.5*,†	0.90 (0.47–1.73)	2.74 (1.29–5.80)*,†
Ever had an STI	3.0	6.3	13.9	2.38 (0.35–16.24)	4.85 (1.25–18.75)*
Positive chlamydia test	1.9	3.3	1.1	1.84 (0.09–37.32)	0.38 (0.03–5.80)
Lack of AIDS knowledge	44.0	47.8	33.9	1.15 (0.73–1.81)	0.66 (0.13–3.30)

\*Significantly different from "none" at  $p < .05$ . †Significantly different from "hit" at  $p < .05$ . ‡Among sexually active respondents (1,554 women and 1,590 men). Note: All multivariate equations included 10-year age-group, urban residence and geographic region; some equations included additional measures—see text for details.