

Table 1. Percentage of U.S. women of reproductive age who reported having engaged in selected pregnancy-related behaviors, by survey, according to the intention status of the birth

Behavior	Survey	Total	Intended	Mistimed	Unwanted	N
Recognized pregnancy in first 6 weeks	NMIHS	67.7	73.2 ^{*,**}	61.1 ^{**}	56.1	9,122
	NSFG	55.9	63.1 ^{*,**}	45.8	43.2	2,535
Made prenatal care visit in first 8 weeks of pregnancy	NMIHS	60.1	67.2 ^{*,**}	51.1	48.3	9,055
	NSFG	40.8	45.1 ^{*,**}	36.0	30.7	2,547
Made recommended no. of prenatal care visits†	NMIHS	67.3	69.4 ^{*,**}	64.9	62.3	8,840
	NSFG	76.5	78.2 [*]	73.2	75.2	2,548
Quit smoking‡	NMIHS	26.4	30.4 ^{*,**}	23.3	19.6	2,732
	NSFG	15.4	16.9 ^{**}	17.8 ^{**}	7.9	496
Reduced or quit alcohol use§	NMIHS	94.0	94.8 ^{**}	94.0 ^{**}	85.7	3,460
	NSFG	na	na	na	na	na
Took vitamins††	NMIHS	91.1	92.5 ^{*,**}	89.7 ^{**}	87.0	8,524
	NSFG	na	na	na	na	na
Gained weight within 5 lbs. of advice received‡‡	NMIHS	41.9	43.9 ^{*,**}	39.7	35.4	6,199
	NSFG	na	na	na	na	na

*Proportion is significantly different from proportion mistimed at $p < .05$. **Proportion is significantly different from proportion unwanted at $p < .05$. †Among those who received any prenatal care. ‡Among smokers only. §Among those who drank prior to pregnancy. ††Among those who were advised to take vitamins. ‡‡Among those who received prenatal care advice on weight gain. *Notes:* All percentages shown here are weighted. NMIHS=National Maternal and Infant Health Survey. NSFG=National Survey of Family Growth. na=not applicable, because not measured in NSFG.