

Table 1. Population estimates (in 000s) of number of U.S. men and women with a self-reported STD in past year and in his or her lifetime (and 95% confidence intervals), by type of STD, 1992 National Health and Social Life Survey

| Disease | N | Total | Men | Women |
|--------------------------|------------|-------------------------------|------------------------------|------------------------------|
| STDs IN PAST YEAR | | | | |
| All | 53 | 2,102 (1,294–2,911) | 1,150 (491–1,809) | 952 (525–1,378) |
| Bacterial† | 29 | 1,229 (658–1,801) | 685 (220–1,150) | 555 (237–853) |
| Viral‡ | 21 | 884 (364–1,404) | 465 (6–924) | 419 (174–664) |
| Gonorrhea | 7 | 304 (47–561) | 156 (0–373) | 148 (101–286) |
| Genital warts | 17 | 750 (240–1,260) | 465 (6–924) | 284 (63–505) |
| Chlamydia | 17 | 748 (288–1,209) | 398 (18–778) | 350 (122–578) |
| STDs IN LIFETIME | | | | |
| All | 614 | 21,788 (18,214–25,363) | 11,585 (9,247–13,923) | 10,203 (8,508–11,899) |
| Bacterial† | 399 | 13,381 (10,950–15,812) | 8,160 (6,324–9,995) | 5,221 (4,335–6,107) |
| Viral‡ | 215 | 10,769 (8,583–12,954) | 4,690 (3,418–5,961) | 6,079 (4,721–7,437) |
| Gonorrhea | 242 | 7,555 (5,901–9,209) | 5,706 (4,250–7,164) | 1,848 (1,418–2,278) |
| Genital warts | 159 | 6,149 (4,634–7,664) | 2,450 (1,467–3,434) | 3,698 (2,705–4,691) |
| Chlamydia | 115 | 3,846 (2,766–4,927) | 1,494 (772–2,216) | 2,352 (1,666–3,038) |
| Genital herpes | 72 | 2,360 (1,525–3,194) | 960 (476–1,443) | 1,400 (887–1,914) |
| Hepatitis | 68 | 2,893 (1,941–3,845) | 1,635 (1,072–2,196) | 1,258 (608–1,908) |
| Nongonococcal urethritis | 33 | 1,616 (1,037–2,195) | 1,616 (1,037–2,195) | na |
| PID | 57 | 1,477 (947–2,008) | na | 1,477 (947–2,008) |
| Syphilis | 27 | 280 (23–537) | 147 (0–330) | 133 (0–300) |
| HIV | 7 | 280 (23–537) | 147 (0–330) | 133 (0–300) |

†Gonorrhea, chlamydia, nongonococcal urethritis, PID and syphilis. ‡Genital warts, herpes, hepatitis and HIV. *Note:* For estimates of STDs in the past year, certain STDs are omitted, as the estimates are unreliable because of small Ns—genital herpes (3), nongonococcal urethritis (3), PID (2), syphilis (1) and HIV (1). The sum of individual estimates of bacterial and viral STDs is slightly larger than the total shown because of rounding. na=not applicable.