

**Table 2. Percentage distribution of respondents who had ever had an STD, by where they said they had gone to receive treatment for the STD, according to age and STD reported**

Disease	N†	Private MD/ group practice	Emergency room	Family Planning clinic	STD clinic	Other clinic	Multiple sites	Site not mentioned	Total
<b>Total</b>	<b>614</b>	<b>48.6 (43.3–53.9)</b>	<b>7.2 (4.5–9.9)</b>	<b>5 (3.0–9.2)</b>	<b>5.4 (3.4–7.4)</b>	<b>7.6 (5.4–9.8)</b>	<b>2.8 (1.8–4.6)</b>	<b>23.4 (22.4–24.4)</b>	<b>100.0</b>
<b>STD</b>									
Gonorrhea	242	26.1 (21.4–31.4)	10.7 (5.8–15.6)	6.7 (3.7–10.4)	10.3 (5.6–15.0)	14.7 (9.4–20.0)	3.3 (0.8–5.8)	28.3 (21.0–35.6)	100.0
Syphilis	28	28.9 (7.3–50.5)	13.2 (1.0–25.4)	8.7 (0.0–24.6)	8.9 (0.0–20.1)	22.7 (8.2–37.2)	0.0	17.7 (2.6–32.8)	100.0
Genital herpes	72	48.9 (36.3–61.4)	1.6 (0.0–4.7)	2.9 (0.0–6.6)	6.4 (0.0–15.6)	4.2 (0.0–8.7)	6.5 (0.2–12.8)	29.5 (20.3–38.7)	100.0
Chlamydia	115	46.8 (36.2–57.4)	4.1 (0.0–9.4)	10.1 (3.0–17.2)	8.7 (2.8–14.6)	2.9 (0.4–5.4)	2.1 (0.0–4.5)	25.3 (14.4–30.8)	100.0
Genital warts	159	58.5 (49.7–67.3)	1.0 (0.0–2.6)	5.0 (0.3–9.7)	5.3 (1.4–9.2)	6.0 (1.7–10.3)	1.5 (0.0–3.7)	22.6 (14.4–30.8)	100.0
Hepatitis	68	61.5 (45.8–67.3)	11.0 (1.0–21.0)	0.0 (0.0)	0.0 (0.0)	4.8 (0.0–10.1)	0.5 (0.0–1.5)	22.3 (20.3–34.6)	100.0
PID‡	57	49.0 (32.9–65.1)	12.7 (1.9–23.5)	2.5 (0–6.2)	0.0 (0.0)	2.5 (0.0–6.2)	0.0	33.4 (15.0–51.8)	100.0
Nongonococcal urethritis§	33	31.1 (14.0–48.2)	4.8 (0.0–11.9)	0.0(0.0)	5.5 (0.0–12.8)	17.5 (4.4–30.6)	3.3 (0.0–9.8)	37.8 (21.5–54.1)	100.0
HIV	7	57.1 (0.0–135.7)	14.3 (0.0–39.8)	0.0(0.0)	14.3 (0.0–39.8)	14.3 (0.0–39.8)	0.0	0.0	100.0

†Total number of each STD reported. ‡Female respondents only. §Male respondents only. *Notes:* All percentages are weighted. Confidence intervals are shown in parentheses.