

**Table 3. Percentage of women discontinuing contraceptive use for method-related reasons (and 95% confidence interval), by method, according to duration of use**

Method	N	Duration of use			
		6 months	12 months	18 months	24 months
<b>Total</b>	<b>6,867</b>	<b>30.6 (29.4–31.8)</b>	<b>43.6 (42.1–45.1)</b>	<b>53.9 (52.2–55.7)</b>	<b>61.3 (59.2–63.4)</b>
Implant	146	6.2 (3.2–11.6)	15.7 (9.9–24.0)	18.3 (11.2–28.4)	20.1 (10.9–34.1)
Pill	2,130	18.0 (16.3–19.8)	32.0 (29.6–34.5)	43.0 (40.0–45.9)	51.3 (47.7–54.8)
IUD	59	18.1 (9.9–30.7)	36.4 (24.0–50.8)	47.8 (31.7–64.4)	59.2 (41.3–75.0)
Diaphragm	166	28.4 (21.5–36.4)	42.8 (33.6–52.6)	53.4 (43.0–63.5)	59.9 (47.3–71.2)
Injectable	209	23.3 (17.2–30.8)	44.4 (34.3–55.1)	54.1 (38.8–68.7)	59.5 (39.0–77.2)
Male condom	2,925	33.9 (31.9–35.9)	47.3 (44.9–49.7)	58.2 (55.4–60.9)	65.8 (62.5–69.1)
Periodic abstinence	250	38.2 (31.9–44.9)	48.8 (41.4–56.3)	60.5 (52.3–68.2)	66.7 (56.0–76.0)
Withdrawal	440	47.4 (42.4–52.5)	57.1 (51.4–62.6)	66.3 (60.3–71.9)	73.4 (66.4–79.3)
Spermicide	164	47.0 (38.7–55.4)	58.2 (48.8–67.1)	65.1 (53.0–75.4)	71.3 (58.3–81.6)
Sponge	111	51.3 (41.1–61.3)	63.7 (51.8–74.1)	85.5 (72.0–93.1)	90.1 (69.3–97.3)
Other	267	89.8 (83.2–94.1)	92.1 (85.0–96.0)	92.1 (85.0–96.0)	92.3 (84.0–96.5)