

Table 5. Percentage of women resuming use of contraceptives after discontinuation, by duration of use (in months), according to method discontinued and subsequent method used

Discontinued method and duration	All	Subsequent method			
		Pill	Sterilization	Male condom	Other
All (N 7,357)					
1	68.1 (67.5–68.7)	17.3	5.4	25.0	20.3
3	75.9 (74.8–77.0)	19.3	5.8	28.9	21.9
6	79.2 (77.9–80.4)	20.4	6.0	30.2	22.6
12	82.0 (80.5–83.4)	21.1	6.1	31.5	23.3
Pill (N=2,568)					
1	65.4 (64.3–66.5)	14.6	5.6	28.0	17.2
3	73.1 (71.1–75.0)	18.4	6.0	30.2	18.6
6	76.2 (73.9–78.3)	20.0	6.2	31.1	19.0
12	79.2 (76.6–81.6)	21.2	6.3	32.1	19.6
Male condom (N=3,046)					
1	67.9 (66.9–68.8)	21.2	4.6	24.0	18.1
3	76.6 (74.8–78.2)	22.4	4.9	30.2	19.1
6	79.8 (77.8–81.7)	23.2	5.1	32.1	19.5
12	82.9 (80.5–85.0)	23.7	5.1	34.1	20.0
Withdrawal (N=443)					
1	72.4 (70.2–74.6)	14.8	2.5	28.5	26.6
3	79.2 (74.1–83.5)	15.3	2.6	32.0	29.3
6	84.7 (78.7–89.3)	15.8	3.0	33.5	32.3
12	86.7 (79.1–91.9)	16.4	3.1	33.9	33.3
Other (N=1,300)					
1	72.1 (70.8–73.4)	14.3	8.1	20.3	29.4
3	78.9 (76.2–81.3)	15.3	8.4	22.4	32.7
6	81.9 (79.1–84.5)	15.8	8.6	23.3	34.2
12	84.0 (80.6–86.8)	16.5	8.7	24.1	34.7