

TABLE 7. Percentage of female adolescents aged 15–19 who knew at least one way of avoiding AIDS, by country, according to marital status

Country	Married	All unmarried	Unmarried and sexually active
South and East Africa			
Ethiopia	68.3	64.4	87.9
Kenya	78.1	80.8	85.4
Malawi	91.3	91.5	93.9
Mozambique	59.3	71.9***	79.1
Namibia	86.6	87.2	90.7
Rwanda	93.6	93.0	96.6
South Africa	97.0	96.0	97.8
Tanzania	81.2	84.0	92.6
Uganda	82.1	85.7	89.4
Zambia	78.8	78.3	80.6
Zimbabwe	74.4	80.8*	82.2
West Africa			
Benin	26.8	22.8	82.3
Burkina Faso	56.7	62.8*	78.7
Cameroon	75.9	84.7***	89.4
Côte d'Ivoire	91.5	95.6	97.5
Gabon	78.9	83.6†	88.0
Ghana	59.1	78.9***	81.2
Guinea	80.5	80.9	84.8
Mali	48.6	56.6**	72.4
Niger	61.6	67.2	67.2
Nigeria	57.3	59.6	64.0
Togo	61.6	74.0***	79.7
Latin America/Caribbean			
Brazil	82.5	90.4***	94.7
Colombia	78.4	89.5***	91.3
Dom. Republic	90.4	91.6	95.6
Guatemala	u	u	u
Haiti	50.4	53.8	68.9
Nicaragua	33.8	40.7**	36.9
Peru	70.4	83.9**	78.2

* $p < .05$. ** $p < .01$. *** $p < .001$. † $p < .10$. Notes: Significance testing is for the difference between married adolescents and all unmarried adolescents. Data from Cameroon, Tanzania and Mozambique come from the DHS surveys prior to those listed in Table 1, because the question about avoiding AIDS was not asked in these countries' most recent surveys. u=unavailable.