

TABLE 3. Selected measures of attitudes toward emergency contraceptive pills, according to country and type of health care provider

Measure	Jamaica					Barbados				
	Total	Nurse	GP	Ob/gyn	Pharmacist	Total	Nurse	GP	Ob/gyn	Pharmacist
Method should be available without prescription	57.0***	63.3	51.1	81.8	50.7	23.2***	0.0	17.9	33.3	40.3
Method encourages sexual risk-taking†	59.1***	76.7	48.4	16.1	84.5	55.1***	61.1	47.6	16.7	63.5
Availability leads to increased STI transmission†	59.3***	46.2	51.1	19.4	92.8	45.9***	52.8	32.9	20.0	59.2
Method is necessary to reduce unintended pregnancies†	88.3**	80.0	90.1	96.9	85.7	75.3*	61.8	79.8	83.3	75.7
Willing to provide method in advance	43.6***	6.7	56.5	78.1	26.8	7.0**	8.3	8.3	16.7	4.1
Willing to provide method to										
Rape victims	90.2***	90.0	92.5	100.0	83.1	83.8***	94.3	94.0	100.0	65.8
Women who experienced condom failure	97.8*	93.1	97.9	100.0	98.6	88.5***	91.7	95.2	100.0	78.4
Women who did not use any contraception	92.9***	80.0	93.5	100.0	94.3	70.0***	58.3	81.0	100.0	60.8
Minors without parental consent	65.9***	69.0	69.7	87.1	49.2	29.7***	25.0	34.9	83.3	21.6
Men requesting method for their partners	50.9***	31.0	55.4	71.9	43.7	28.5***	13.9	15.5	33.3	50.0
Any woman regardless of circumstances	25.8***	27.6	28.9	43.8	12.9	18.2**	8.6	18.1	50.0	20.3

*p<.05. **p<.01. ***p<.001. †Includes respondents who "somewhat" or "totally" agreed with these statements; respondents who answered "don't know" are excluded. Notes: Missing data are excluded. Statistical comparisons are across provider groups within each country. GP=general practitioner. Ob/gyn=obstetrician-gynecologist.