

**TABLE 3. Odds ratios (and 95% confidence intervals) from logistic regression analyses predicting the effects of various characteristics and perceptions on the likelihood of condom use at last intercourse with a main partner, by sex**

Variable	Females	Males
<b>Characteristic</b>		
Used hormonal contraceptive at last intercourse	0.42 (0.24–0.74)**	0.26 (0.09–0.77)*
STD clinic	1.41 (0.78–2.54)	1.66 (0.59–4.66)
Age	0.74 (0.60–0.92)**	0.74 (0.50–1.09)
Relationship length	1.00 (0.98–1.01)	0.98 (0.95–1.01)
$\chi^2$	24.1**	20.8**
<b>Perception</b>		
Negative view of becoming pregnant	1.08 (0.99–1.18)	1.13 (0.96–1.33)
At risk of becoming pregnant	1.06 (0.93–1.21)	0.99 (0.75–1.29)
Negative view of acquiring STD	1.08 (1.00–1.16)	1.16 (1.00–1.34)*
At risk of acquiring STD	1.06 (1.02–1.11)**	1.07 (0.99–1.17)
$\chi^2$	20.0**	13.5**
<b>Two-way interactions</b>		
Hormonal contraceptive use x pregnancy negative	0.75 (0.62–0.91)**	0.82 (0.56–1.19)
Hormonal contraceptive use x at risk of acquiring STD	1.17 (1.05–1.30)**	1.05 (0.82–1.33)
Pregnancy negative x at risk of acquiring STD	1.00 (0.98–1.01)	0.97 (0.94–1.00)†
$\chi^2$	17.8**	4.1
<b>Three-way interaction</b>		
Hormonal contraceptive use x pregnancy negative x at risk of acquiring STD	0.95 (0.91–0.99)*	0.82 (0.69–0.99)*
$\chi^2$	6.3*	7.5**
$\chi^2$ for model	68.2**	45.9**

\*p<.05. \*\*p<.01. †p<.10.