

TABLE 4. Odds ratios (and 95% confidence intervals) from logistic regression analyses predicting the likelihood of female adolescents' condom use at last intercourse with a main partner, by selected characteristics, according to adolescents' perceptions about becoming pregnant and their risk of acquiring an STD

| Variable | Negative view of becoming pregnant | | Risk of acquiring STD | |
|-------------------------------------------------------|------------------------------------|------------------|-----------------------|--------------------|
| | More (N=128) | Less (N=146) | Higher (N=131) | Lower (N=143) |
| Used hormonal contraceptive at last intercourse | 0.14 (0.06–0.36)** | 0.79 (0.39–1.61) | 0.86 (0.38–1.94) | 0.21 (0.09–0.46)** |
| STD clinic | 3.55(1.37–9.23)** | 1.02 (0.49–2.14) | 1.70 (0.77–3.75) | 1.58 (0.70–3.55) |
| Age | 0.68 (0.48–0.95)* | 0.77 (0.58–1.01) | 0.81 (0.61–1.07) | 0.74 (0.54–1.02) |
| Relationship length | 1.00 (0.97–1.03) | 1.00 (0.98–1.02) | 1.00 (0.98–1.02) | 0.98 (0.95–1.00) |
| χ^2 | 29.5** | 5.5 | 6.0 | 26.1** |

*p<.05. **p<.01.