

TABLE 1. Percentage of married women using selected contraceptive methods who switched to a new method within two years, by new (destination) method, 1995 National Survey of Family Growth

Origin method	N	Destination method							
		All	Female sterilization	Male sterilization	Implant/injectable/IUD	Pill	Condom	Other reversible	None
All	2,973	39.6 (3.7–12.5)	2.0 (1.3–2.8)	2.8 (2.1–3.7)	2.4 (1.6–3.2)	7.9 (6.5–9.2)	9.3 (7.8–10.9)	6.6 (5.3–7.9)	8.7 (7.2–10.4)
Implant/injectable/IUD	147	30.0 (20.1–40.2)	4.0 (0.2–11.2)	0.0 (na)	na	5.1 (1.4–9.6)	9.2 (3.6–16.3)	6.1 (1.9–11.9)	5.5 (1.4–10.3)
Pill	745	41.5 (36.0–47.1)	2.2 (0.8–3.9)	1.7 (0.6–3.0)	3.3 (1.5–5.3)	na	14.7 (11.1–19.0)	9.0 (6.0–12.2)	10.6 (7.3–14.6)
Condom	764	41.6 (36.0–47.2)	1.5 (0.6–2.7)	5.6 (3.3–8.1)	2.6 (1.2–4.3)	11.5 (8.2–15.3)	na	7.7 (5.4–10.4)	12.6 (9.3–16.1)
Other reversible	546	29.6 (24.3–35.1)	1.5 (0.2–3.3)	2.9 (1.5–4.6)	0.9 (0.1–2.0)	5.7 (3.6–8.2)	8.4 (5.8–11.3)	na	10.2 (6.7–14.1)
None	771	42.8 (37.4–48.5)	2.1 (1.0–3.4)	1.5 (0.5–2.8)	2.3 (1.2–3.8)	17.0 (12.7–21.9)	12.4 (9.5–15.7)	7.4 (4.7–10.6)	na

Notes: Ns are unweighted. Figures in parentheses are 95% confidence intervals, derived from bootstrapping using 1,000 resamples. na=not applicable.