

**TABLE 5. Hazard model coefficients indicating the effects of married women's characteristics on the risk of switching from a specific origin method to a specific destination method**

Origin method and characteristic	Destination method						
	All	Sterilization	Implant/ injectable/IUD	Pill	Condom	Other reversible	None
<b>Pill</b>							
Age at start of interval	0.004	0.039*	-0.016	na	-0.016	0.016	-0.091*
Black	0.238	0.776*	-0.524	na	0.265	-0.205	0.322*
Hispanic	0.190	-1.140*	0.181	na	0.013	-0.312	0.126
Completed yrs. of education	-0.028	-0.021	-0.031	na	0.175*	-0.113*	-0.070*
Parity 0 at start of interval	0.461	-0.888*	1.499*	na	-0.035	-0.211	0.173
No religious affiliation	0.214	0.504*	1.491*	na	-0.308	0.000	-1.583*
<b>Condom</b>							
Age at start of interval	-0.024	0.041*	-0.035*	-0.003	na	-0.187*	-0.019
Black	-0.010	0.531*	-0.187	-1.386*	na	-0.900*	0.235
Hispanic	-0.273	-2.272*	-0.248	-2.217*	na	-0.842*	-0.236
Completed yrs. of education	-0.021	-0.136*	-0.153*	-0.119*	na	0.008	0.043
Parity 0 at start of interval	0.014	†	†	0.061	na	-1.034*	0.263
No religious affiliation	0.103	-2.047*	-0.885	0.155	na	-0.187	-0.283
<b>Other reversible</b>							
Age at start of interval	-0.019	0.056*	0.064*	-0.121*	-0.089*	na	0.075*
Black	0.229	-1.825*	0.328	0.217	-1.738*	na	-0.798*
Hispanic	0.176	-1.977*	1.114*	-0.089	0.520*	na	-0.160
Completed yrs. of education	0.022	0.109*	0.066*	0.161*	0.043	na	-0.133*
Parity 0 at start of interval	0.425	†	†	1.373*	0.258	na	1.694*
No religious affiliation	0.117	0.875*	†	-0.021	0.629*	na	0.418*
<b>None</b>							
Age at start of interval	-0.036*	0.091*	-0.042	-0.081*	-0.027	0.036	na
Black	0.056	0.884*	0.055	0.821*	0.436*	-1.661*	na
Hispanic	0.612*	1.012*	1.636*	0.127	0.763*	-0.001	na
Completed yrs. of education	0.086*	0.090*	-0.074	0.090*	0.062	0.211*	na
Parity 0 at start of interval	-0.700*	-1.887*	†	-0.547*	0.126	-2.010*	na
No religious affiliation	0.148	-1.113	0.900	-1.725*	1.236*	-0.438	na

\*p<.05. †Variable deleted from the model because no women with the characteristic switched to the method. Notes: Models also control for women's living arrangements during childhood, age at menarche and duration of use (not shown). na=not applicable. No results are shown for models in which long-term reversible methods were the origin method because few married women use those methods and rates of switching from those methods to each destination method are very low.