

TABLE 2. Adjusted odds ratios (and 95% confidence intervals) from logistic regression analyses assessing the effects of selected characteristics on women's belief that they never have certain sexual rights

| Characteristic | Odds ratio | Characteristic | Odds ratio |
|--|------------------|---|------------------|
| To make own decisions about sexual activity, regardless of partner's wishes | | To masturbate to orgasm (cont'd.) | |
| 1–2 lifetime partners* | 2.48 (1.33–4.62) | 1–2 lifetime partners* | 1.66 (1.15–2.40) |
| History of sexual assault | 1.96 (1.07–3.57) | ≥1 birth | 1.59 (1.17–2.16) |
| To make own decision about birth control, regardless of partner's wishes | | To tell partner he is being too rough | |
| 1–2 lifetime partners* | 1.96 (1.01–2.57) | Black† | 1.71 (1.08–2.70) |
| To tell partner "I want to make love" | | Hispanic† | 1.90 (1.19–3.05) |
| Black† | 2.14 (1.10–4.16) | 1–2 lifetime partners* | 1.81 (1.14–2.89) |
| Hispanic† | 2.04 (1.07–4.04) | No history of physical assault | 1.60 (1.05–2.41) |
| 1–2 lifetime partners* | 3.26 (1.64–6.68) | To tell partner "I want to be hugged or cuddled without sex" | |
| To tell partner "I do not want to make love" | | Black† | 3.09 (1.46–6.54) |
| Black† | 2.18 (1.12–4.25) | Hispanic† | 2.58 (1.20–5.63) |
| C average‡ | 2.09 (1.18–3.72) | <C average‡ | 2.06 (1.01–4.23) |
| <C average‡ | 2.81 (1.28–6.14) | 1–2 lifetime partners* | 2.85 (1.36–5.99) |
| 1–2 lifetime partners* | 2.63 (1.31–5.26) | Inconsistent contraceptive use | 1.85 (1.10–3.12) |
| Inconsistent contraceptive use | 1.72 (1.03–2.87) | To tell relative "I am not comfortable being hugged or kissed in certain ways" | |
| No history of physical assault | 2.05 (1.08–3.89) | Hispanic† | 1.58 (1.07–2.33) |
| To tell partner "I won't have sex without birth control" | | To ask partner if he has been examined for STDs | |
| Black† | 1.97 (1.22–3.17) | 18–21§ | 1.66 (1.02–2.71) |
| Hispanic† | 1.82 (1.10–2.99) | Hispanic† | 1.67 (1.02–2.72) |
| <C average‡ | 1.86 (1.06–3.27) | Inconsistent contraceptive use | 1.50 (1.02–2.21) |
| Inconsistent contraceptive use | 1.93 (1.32–2.81) | To stop foreplay at any time, including at the point of intercourse | |
| To tell partner "I want to make love differently" | | Black† | 1.75 (1.10–2.79) |
| 14–17§ | 2.25 (1.31–2.87) | Hispanic† | 1.65 (1.02–2.68) |
| Black† | 2.45 (1.48–4.07) | <C average‡ | 1.75 (1.01–3.07) |
| Hispanic† | 2.58 (1.54–4.25) | 1–2 lifetime partners* | 1.98 (1.22–3.21) |
| <C average‡ | 1.85 (1.05–3.27) | Inconsistent contraceptive use | 1.49 (1.04–2.14) |
| 1–2 lifetime partners* | 2.46 (1.50–4.03) | ≥1 birth | 1.66 (1.13–2.45) |
| To masturbate to orgasm | | To refuse to have sex even if she has enjoyed it with this partner before | |
| 14–17§ | 2.06 (1.37–3.11) | Hispanic† | 1.91 (1.18–3.07) |
| 18–21§ | 1.43 (1.01–2.01) | <C average‡ | 1.81 (1.03–3.19) |
| Black† | 1.69 (1.20–2.37) | 1–2 lifetime partners* | 1.88 (1.16–3.04) |
| Hispanic† | 2.48 (1.73–3.57) | Inconsistent contraceptive use | 1.52 (1.05–2.19) |
| C average‡ | 1.36 (1.01–1.85) | No history of physical assault | 1.65 (1.08–2.53) |
| <C average‡ | 2.34 (1.43–3.83) | | |

*Reference group is six or more lifetime partners. †Reference group is white. ‡Reference group is B average or higher. §Reference group is 22–26-year-olds. Notes: Consistency of contraceptive use reflects clients' use in the past year. Results are presented only for associations that are significant at $p < .05$.