

TABLE 1. Percentage distribution of women obtaining abortions in 2000, by contraceptive method used in the month of conception, and of women at risk of unintended pregnancy in 1995, by contraceptive method used

Method	Women having abortions, 2000	Women at risk of unintended pregnancy, 1995†
Any method	53.7	92.5
Long-acting	1.1	40.5
Sterilization	0.1	35.8
IUD	0.1	0.7
Implant/injectable	0.9	4.0
Pill	13.6	24.9
Male condom	27.6	18.9
Withdrawal	7.3	2.8
Periodic abstinence	2.2	2.1
Other‡	1.9	3.3
No method	46.3	7.5
Never used	8.1	0.1
Previously used	38.2§	7.4
Total	100.0	100.0
Unweighted N	10,683	7,725
Weighted N (in 000s)	1,313	41,796

†Based on special tabulations from the 1995 National Survey of Family Growth. Includes women using a contraceptive method in the survey month and fertile women using no method who had intercourse in the previous three months and were not pregnant, seeking pregnancy or postpartum and whose partner was not sterile. ‡Female condom, diaphragm, foam, sponge, suppository or any other method. §Includes women who used only emergency contraceptive pills to prevent the current pregnancy. *Source:* Weighted number of women having abortions from reference 5.