

TABLE 6. Percentages of pill and condom users obtaining abortions, by perceived reasons for pregnancy

Reason	%
PILL USERS	(N=1,458)
Perfect use/method failed	12.7
Inconsistent use	75.9
Forgot to take pills	45.1
Away from home and didn't have pills	15.9
Ran out of supplies	10.3
Sick	7.7
Didn't think I would have sex again	2.1
Didn't feel like taking pills	1.6
Partner didn't want me to take	0.3
Thought I might want to get pregnant	0.2
Other reason	3.6
Other†	14.9
CONDOM USERS	(N=2,945)
Perfect use/method failed	13.5
Inconsistent use	49.3
Didn't think I would get pregnant	20.4
Didn't have it with me	14.3
Not expecting to have sex	12.8
Forgot	7.1
Didn't feel like using	5.6
Partner didn't feel like using	3.5
Partner was supposed to bring	1.0
Partner wanted me to get pregnant	1.0
I was forced to have sex	0.5
Thought I might want to get pregnant	0.5
Didn't care if I got pregnant	0.3
Other reason	1.8
Condom broke/slipped	41.6
Other‡	3.2

†Included use with other drugs or antibiotics, absorption problems (vomiting or diarrhea), change of pill or dose, and not finishing first pack. ‡Included incorrect use and use of lambskin condoms and expired condoms. *Note:* Users could indicate more than one reason for imperfect use.