

**TABLE 3. Adjusted odds ratios (and 95% confidence intervals) from multinomial logistic regression analyses assessing variables associated with partially inadequate or inadequate prenatal care versus adequate care**

Variable	Partially inadequate	Inadequate
<b>LEVEL I</b>		
<b>Age</b>	0.99 (0.96–1.02)	0.99 (0.96–1.03)
<b>Income</b>		
1st tertile (ref)	1.00	1.00
2nd tertile	0.69 (0.43–1.13)	0.57 (0.35–0.94)***
3rd tertile	0.43 (0.26–0.72)***	0.14 (0.07–0.28)***
<b>Education</b>	0.96 (0.88–1.04)	0.92 (0.85–1.01)
<b>Skin color</b>		
White (ref)	1.00	1.00
Nonwhite	1.06 (0.66–1.72)	1.26 (0.77–2.07)
<b>LEVEL II</b>		
<b>Parity</b>		
1 (ref)	1.00	1.00
2–3	1.81 (1.07–3.06)***	3.93 (2.05–7.53)***
≥4	3.30 (1.58–6.91)***	9.03 (3.82–16.30)***
<b>Living with child's father</b>		
Yes (ref)	1.00	1.00
No	0.76 (0.38–1.50)	2.74 (1.53–4.92)***
<b>LEVEL III</b>		
<b>Planned pregnancy</b>		
Yes (ref)	1.00	1.00
No	1.34 (0.82–2.20)	1.96 (1.01–3.79)*
<b>Satisfied with pregnancy (prenatal)</b>		
Yes (ref)	1.00	1.00
No	1.19 (0.69–2.04)	2.12 (1.22–3.67)**
<b>Attempted abortion</b>		
Yes	1.37 (0.42–4.55)	1.55 (0.51–4.68)
No (ref)	1.00	1.00

\*p<.05. \*\*p<.01. \*\*\*p≤.001. Notes: Estimates are adjusted for the hospital where delivery occurred and maternal age. Variables that were not significant at p<.05 for either outcome were not included in the final model. ref=reference group.