

TABLE 4. Number of women seeking postabortion care; number and percentage distribution of these women by severity of morbidity; and percentage of these women who were hospitalized—all according to region

Region	All		Severe		Moderate		Low		% hospitalized
	No.	No.	%	No.	%	No.	%		
Addis Ababa	7,233	868	12***	651	9	5,714	79	13 (9–18)**	
Other	50,731	14,816	29	7,052	14	28,863	57	24 (20–27)	
Dire Dawa/Harari	3,671	918	25	220	6	2,533	69	31 (15–48)	
SNNP	10,896	2,724	25	1,416	13	6,756	62	20 (13–27)	
Oromiya	20,310	6,093	30	2,843	14	11,374	56	26 (19–33)	
Amhara	9,242	2,865	31	1,386	15	4,991	54	14 (8–19)	
ASBG	3,274	1,048	32	786	24	1,440	44	24 (16–33)	
Tigray	3,338	1,168	35	401	12	1,769	53	46 (34–58)	

Percentage for Addis Ababa is significantly different from percentage for other regions at $p < .01$. *Percentage distribution by severity of morbidity in Addis Ababa is significantly different from that in other regions at $p < .001$. Notes: All data are weighted. Hospitalization refers to a stay of at least 24 hours. SNNP= Southern Nations, Nationalities and People's Region. ASBG= Affar, Somali, Benishangul-Gumuz and Gambela.