

TABLE 1. Selected characteristics of pelvic pain conditions among women aged 25–54 in Hermosillo, Mexico, 2005

Characteristic	Menstrual pain†	Pain with intercourse‡	Chronic pelvic pain§
ALL WOMEN	(N=1,007)	(N=1,183)	(N=1,201)
With condition (%)	39.9 (36.9–43.0)	12.3 (10.5–14.4)	6.4 (5.1–8.0)
Pain severity (%)††			
Slight	15.5 (13.4–17.9)	6.7 (5.4–8.3)	2.4 (1.7–3.5)
Moderate	16.4 (14.2–18.9)	4.6 (3.6–6.1)	3.2 (2.3–4.4)
Severe	7.9 (6.3–9.7)	1.0 (0.6–1.8)	0.7 (0.3–1.4)
WOMEN WITH CONDITION	(N=402)	(N=146)	(N=77)
Age at onset (%)‡‡			
≤24 yrs.	72.2	29.0	na
≥25 yrs.	27.7	71.0	na
Mean yrs. since first symptoms‡‡	15.8 (SD=10.2)	5.8 (SD=6.3)	na

†Among women who had had at least one menstrual period in the last 12 months. ‡Among women who had had at least one sexual partner in the last 12 months. §Among women who had not been pregnant in the previous 12 months. ††Data are missing for two women on menstrual pain and chronic pelvic pain. ‡‡Data are missing for six women on menstrual pain and one woman on pain with intercourse. Notes: Unless otherwise noted, figures in parentheses are 95% confidence intervals. na=not applicable. SD=standard deviation.