

**TABLE 2. Percentage of women having pelvic pain conditions, and the crude odds ratios (and 95% confidence intervals) of having each condition from logistic regression analyses, according to selected characteristics, by condition**

Characteristic	Menstrual pain†			Pain with intercourse‡			Chronic pelvic pain§		
	N	%	OR (95% CI)	N	%	OR (95% CI)	N	%	OR (95% CI)
<b>DEMOGRAPHIC</b>									
<b>Age (yrs.)</b>									
25–34	450	45.3	1.50 (1.16–1.93)	436	17.4	2.04 (1.44–2.89)	397	10.3	2.45 (1.54–3.91)
≥35 (ref)	557	35.5	1.00	747	9.4	1.00	804	4.4	1.00
<b>Marital status</b>									
Single	136	39.7	1.01 (0.69–1.47)	88	5.6	0.41 (0.1–1.04)	142	3.5	0.51 (0.20–1.31)
Married/living as married (ref)	792	39.4	1.00	1,026	12.6	1.00	941	6.6	1.00
Divorced/separated/widowed	79	45.5	1.28 (0.81–2.05)	69	15.9	1.30 (0.66–2.55)	118	8.4	1.31 (0.65–2.63)
<b>REPRODUCTIVE</b>									
<b>Age at sexual debut (yrs.)</b>									
10–14	42	57.1	2.26 (1.20–4.26)	51	23.5	2.54 (1.28–5.02)	46	6.5	1.16 (0.35–3.09)
15–17	239	46.0	1.45 (1.07–1.47)	279	15.1	1.46 (0.98–2.17)	278	9.0	1.65 (0.99–2.74)
≥18 (ref)	686	37.0	1.00	852	10.8	1.00	834	5.6	1.00
No sexual debut	40	35.0	0.91 (0.47–1.78)	na	na	na	43	4.7	0.81 (0.19–3.48)
Missing	0			1			0		
<b>Type of birth</b>									
Vaginal (ref)	524	38.3	1.00	41	10.7	1.00	661	6.5	1.00
Cesarean	258	40.7	1.10 (0.81–1.49)	662	15.9	1.58 (1.06–2.35)	272	6.2	0.95 (0.53–1.71)
Both	145	42.0	1.16 (0.80–1.69)	294	11.3	1.06 (0.63–1.78)	182	6.0	0.92 (0.46–1.83)
No births	80	43.7	1.25 (0.77–2.01)	185	17.0	1.71 (0.73–4.00)	86	6.9	1.07 (0.44–2.61)
Missing	0			1			0		
<b>IUD use</b>									
Yes	107	45.8	1.30 (0.87–1.95)	110	20.0	1.91 (1.15–3.16)	103	6.8	1.07 (0.47–2.39)
No (ref)	900	39.2	1.00	1,073	11.5	1.00	1,098	6.3	1.00
<b>History of STI/PID</b>									
Yes	126	57.9	2.31 (1.58–3.38)	153	24.1	2.71 (1.78–4.13)	151	16.5	3.80 (2.28–6.34)
No (ref)	880	37.2	1.00	1,028	10.5	1.00	1,049	4.9	1.00
Missing	1			2			1		
<b>History of uterine fibroids</b>									
Yes	38	57.9	2.13 (1.10–4.11)	70	17.1	1.51 (0.79–2.88)	76	14.4	2.70 (1.36–5.36)
No (ref)	962	39.2	1.00	1,105	12.0	1.00	1,120	5.8	1.00
Missing	7			8			5		
<b>MEDICAL</b>									
<b>History of colitis</b>									
Yes	321	50.4	1.91 (1.46–2.51)	393	17.8	2.08 (1.54–2.97)	402	8.7	1.71 (1.07–2.72)
No (ref)	683	34.7	1.00	787	9.4	1.00	796	5.3	1.00
Missing	3			3			3		
<b>Chronic urinary tract infections</b>									
Yes	380	45.5	1.45 (1.12–1.89)	483	16.9	2.06 (1.45–2.92)	473	9.0	2.38 (1.28–3.24)
No (ref)	626	36.4	1.00	698	9.0	1.00	727	4.6	1.00
Missing	1			2			1		
<b>ANTHROPOMETRIC</b>									
<b>Body mass index (kg/m<sup>2</sup>)</b>									
<25	238	34.0	1.00	245	11.4	1.00	261	6.5	1.00
≥25	757	41.8	1.39 (1.03–1.89)	924	12.7	1.13 (0.73–1.75)	926	6.4	0.96 (0.55–1.68)
Missing	12			14			14		
<b>Height (cm)</b>									
≤150	73	39.7	0.98 (0.60–1.60)	85	11.7	0.93 (0.47–1.84)	92	11.9	2.12 (1.07–4.17)
>150 (ref)	924	40.0	1.00	1,086	12.5	1.00	1,097	6.0	1.00
Missing	10			12			12		

†Among women who had had a menstrual period in the last 12 months. ‡Among women who had had at least one sexual partner in the last 12 months. §Among women who had not been pregnant in the last 12 months. Notes: ref=reference group. na=not applicable.