

**TABLE 2. Percentage of women aged 15–44 using contraceptives, by method, according to marital status, Georgia Reproductive Health Survey, 1999 and 2005**

Method	1999				2005			
	All (N= 7,798)	Married (N= 5,177)	Previously married (N=517)	Never married (N=2,104)	All (N= 6,376)	Married (N= 4,119)	Previously married (N=386)	Never married (N=1,871)
<b>Any</b>	<b>24.7</b>	<b>40.5</b>	<b>2.7</b>	<b>0.0</b>	<b>28.4</b>	<b>47.3*</b>	<b>4.8</b>	<b>0.2</b>
<b>Modern</b>	<b>12.1</b>	<b>19.8</b>	<b>2.7</b>	<b>0.0</b>	<b>16.1</b>	<b>26.6*</b>	<b>4.8</b>	<b>0.2</b>
IUD	5.9	9.7	0.7	0.0	7.0	11.6	0.5	0.0
Condom	3.9	6.3	1.1	0.0	5.3	8.7	2.0	0.2
Pill	0.6	1.0	0.0	0.0	1.9	3.2	1.0	0.0
Female sterilization	1.0	1.6	0.9	0.0	1.4	2.2	0.0	0.0
Spermicide	0.1	0.1	0.0	0.0	0.5	0.9	1.0	0.0
Injectable	0.0	0.0	0.0	0.0	0.0	0.0	0.2	0.0
Other	0.6	1.0	0.0	0.0	0.0	0.0	0.0	0.0
<b>Traditional</b>	<b>12.6</b>	<b>20.7</b>	<b>0.0</b>	<b>0.0</b>	<b>12.3</b>	<b>20.7</b>	<b>0.0</b>	<b>0.0</b>
Withdrawal	6.4	10.5	0.0	0.0	6.6	11.2	0.0	0.0
Periodic abstinence	6.2	10.2	0.0	0.0	5.7	9.5	0.0	0.0
<b>None</b>	<b>75.2</b>	<b>59.5</b>	<b>97.4</b>	<b>100.0</b>	<b>71.6</b>	<b>52.7</b>	<b>95.2</b>	<b>99.8</b>

\*Difference between 1999 and 2005 is significant at  $p < .05$ . Note: Women in consensual unions are categorized as married.