

**TABLE 4. Odds ratios (and 95% confidence intervals) from multivariate analysis assessing characteristics associated with chlamydia infection among women aged 15–24 visiting family planning clinics, by race**

Characteristic	White	Black
<b>Age</b>		
15–17	1.50 (1.23–1.83)	3.03 (2.04–4.48)
18–21	1.72 (1.45–2.04)	2.11 (1.48–2.99)
22–24 (ref)	1.00	1.00
<b>Reason for visit</b>		
STD	1.16 (0.98–1.37)	0.99 (0.73–1.36)
Family planning (ref)	1.00	1.00
<b>Risk history†</b>		
None (ref)	1.00	1.00
Any	2.23 (1.95–2.56)	1.42 (1.05–1.90)
Partner with STD symptoms	4.45 (3.39–5.84)	2.18 (1.35–3.55)
New partner	1.96 (1.68–2.30)	1.33 (0.94–1.88)
Multiple partners	2.76 (2.23–3.43)	1.67 (0.98–2.85)
<b>Clinical signs</b>		
None (ref)	1.00	1.00
Any	2.53 (2.14–3.00)	2.50 (1.80–3.48)
Cervicitis	2.54 (2.07–3.11)	2.76 (1.84–4.13)
PID	2.69 (1.73–4.19)	0.63 (0.15–2.65)
Cervical friability	3.90 (3.09–4.93)	2.88 (1.78–4.68)
Urethritis	1.54 (0.87–2.73)	1.44 (0.50–4.16)
<b>Symptoms</b>		
No (ref)	1.00	1.00
Yes	1.10 (0.87–1.40)	0.91 (0.63–1.31)
<b>Gonorrhea infection</b>		
No (ref)	1.00	1.00
Yes	13.10 (8.7–19.8)	5.43 (3.55–8.32)

†During the previous year. Note: ref=reference group.